

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International BureauReference No.: AJ  
Filing Date: August 19, 2004  
Application No.: 10/505,315

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 15/82, C07K 14/32, 16/18, G01N 33/53, C07K 14/435, C12N 15/16, 15/62, A01H 5/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/44137</b>  <b>(43) International Publication Date:</b> 8 October 1998 (08.10.98)
<b>(21) International Application Number:</b> PCT/EP98/01952 <b>(22) International Filing Date:</b> 2 April 1998 (02.04.98)  <b>(30) Priority Data:</b> 08/838,219 3 April 1997 (03.04.97) US 08/832,263 3 April 1997 (03.04.97) US 08/832,265 3 April 1997 (03.04.97) US  <b>(71) Applicant (for all designated States except AT US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  <b>(71) Applicant (for AT only):</b> NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1235 Vienna (AT).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ESTRUCH, Juan Jose [ES/US]; 4301 Talcott Drive, Durham, NC 27705 (US). YU, Cao-Guo [CA/US]; 9G Estes Park Apartment, 306 Estes Drive Ext., Carrboro, NC 27510 (US). WARREN, Gregory, Wayne [US/US]; 324 Bond Lake Drive, Cary, NC 27513 (US). DESAI, Nalini, Manoj [US/US]; 107 Silverwood Lane, Cary, NC 27511 (US). KOZIEL, Michael, Gene	<b>[US/US];</b> 14840 Woodcrest, Clive, IA 50325 (US). NYE, Gordon, James [US/US]; 6329 Deerview Drive, Raleigh, NC 27606 (US).  <b>(74) Agent:</b> BECKER, Konrad; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, VJ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> PLANT PEST CONTROL		
<b>(57) Abstract</b>  The present invention is drawn to a novel class of proteins, and their receptors. Novel processes, assays and methods for controlling plant pests are provided.  <div style="text-align: center;"> <p>1 20</p> <p>SGSPGLQEFAAASTMYSRIFLLVIVCAVKASLFTVN</p> <p>40 60</p> <p>VYDDNPETEIASSLKGCNPQECQRCRRLKFPGGA</p> <p>80 100</p> <p><u>CVNGRCKCDNFLSVKDDVSVEEPAILKDLVSLAEQ</u></p> <p>120 140</p> <p>AAKSRCRNRVCDVCRALHNTSGACVDGQCKCTN</p> <p>160</p> <p>KISAGDIVSDPAESLRTCNPIRCDEQCRRNGHEFGV</p> <p>180 200</p> <p><u>CFKGQCKCDYFLKEEVDEPEVTSLPKNCNPQECQ</u></p> <p>220 240</p> <p>RCRRLKFPGGACVNGRCKCDNFFSAGDIVSDPAES</p> <p>260 280</p> <p>LRSCNPIRCDEQCRRNGHEFGVCFKGQCKCDYFL</p> <p>300</p> <p>NSEVDVAVNEFPQAGSKRYCNLTQCNQTCANRFYD</p> <p>320 340</p> <p>SARVIHGWCKCYSKMERQDASPLNDVTEDENEVS</p> <p>360 380</p> <p>NDILRTVAEELSDVSPRACKSASCNQACRAFYFKG</p> <p>396</p> <p>GWCRFGRCQCF</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PLANT PEST CONTROL

The present invention relates to a novel class of proteins for the control of plant pests.

Plant pests are a major factor in the loss of the world's commercially important agricultural crops resulting both in economic hardship to farmers and nutritional deprivation for local populations in many parts of the world. Broad spectrum chemical pesticides have been used extensively to control or eradicate pests of agricultural importance. There is, however, substantial interest in developing effective alternative pesticides.

Control of various pests through the use of biological molecules has been possible in only a limited number of cases. The best known examples of biological molecules with pesticidal uses are the  $\delta$ -endotoxins from *Bacillus thuringiensis* (Bt), which is a gram-positive spore forming microorganism. Varieties of Bt are known that produce more than 25 different but related  $\delta$ -endotoxins. Bt strains produce  $\delta$ -endotoxins during sporulation the use of which is limited because they are active against only a very few of the many insect pests.

The limited specificity of the Bt endotoxins is dependent, at least in part, on both the activation of the toxin in the insect gut (Haider, M. Z. *et al.*, 1986, Eur. J. Biochem. 156:531-540) and its ability to bind to specific receptors present on the insect's midgut epithelial cells (Hofmann, C. P. *et al.*, 1988, PNAS 85:7844-7848). Therefore, the ability to control a specific insect pest using  $\delta$ -endotoxins at present depends on the ability to find an appropriate  $\delta$ -endotoxin with the desired range of activity. In many cases, no such  $\delta$ -endotoxin is known, and it is not certain that one even exists.

Plants also routinely become infected by fungi and bacteria, and many microbial species have evolved to utilize the different niches provided by the growing plant. In addition to infection by fungi and bacteria, many plant diseases are caused by nematodes which are soil-borne and infect roots, typically causing serious damage when the same crop species is cultivated for successive years on the same area of ground.

The severity of the destructive process of disease depends on the aggressiveness of the phytopathogen and the response of the host, and one aim of most plant breeding programs is to increase the resistance of host plants to disease. Novel gene sources and combinations developed for resistance to disease have typically only had a limited period of

successful use in many crop-pathogen systems due to the rapid evolution of phytopathogens to overcome resistance genes.

It is apparent, therefore, that scientists must constantly be in search of new methods with which to protect crops against plant pests. It has been found in the present invention a novel class of proteins which can be used to control plant pests.

Programmed cell death is a process whereby developmental or environmental stimuli activate a genetic program that culminate in the death of the cell (Jacobson, M.D. *et al.*, 1997, Cell 88: 347-354). This genetic potential exists in most, if not all, multicellular organisms. In the case of invertebrates, programmed cell death appears to play a dual role by being an integral part of both the insect development process and a response mechanism to infections particularly of viral nature (Clem, R.J. *et al.*, 1991, Science 254: 1388-1390). Programmed cell death appears to be executed in several different manners leading to either apoptosis, atrophy or differentiation. Apoptosis is one of the best characterized types of programmed cell death encompassing cytological changes including membrane-bound apoptotic bodies and cytoplasmic blebbing as well as molecular changes such as endonucleolysis typified by the generation of oligosomal length fragments (Vaux, D.L and Strasser, A., 1996, PNAS 93:2239-2244). Although the overall apoptotic phenomenology is rather conserved among the different organisms, it is interesting to point out that, for many insect cells, cytoplasmic vacuolization and swelling rather than condensation seem to be the cytological features associated with apoptotic processes (Bowen, I.D., *et al.*, 1996, Micros. Res. Techniq. 34:202-217). The novel class of proteins disclosed within the present invention are shown to induce programmed cell death and exert a pesticidal effect.

The present invention is drawn to VIP3A(c) proteins including homologues thereof. Also provided by the invention are domains of proteins of the VIP3 class, including the toxic domain and the stabilizing domain. A preferred embodiment of the invention is the toxic domain of the VIP3A(a) protein and homologues thereof. Another preferred embodiment are antibodies to proteins of the VIP3 class, but preferably to the VIP3A(c) protein.

The invention also provides hybrid toxins comprising a toxic domain of a protein of the VIP3 class. In a preferred embodiment, the hybrid toxin is a chimeric proteins having a toxic core domain operably linked to a heterologous stabilizing domain. In another preferred embodiment, the hybrid toxin comprises an antibody, or immunologically-active



fragment thereof, which immunologically recognizes the VIP3 receptor operably linked to a toxic domain from other proteins, wherein the toxin domain is obtained from a number of cytotoxic proteins including but not limited to *Bacillus* toxins, including endotoxins and vegetative insecticidal proteins.

Also encompassed by the invention are plants comprising a DNA sequence which encodes a protein of the VIP3 class, but preferably a VIP3A(c) protein. Preferred embodiments include plants selected from the group consisting of maize, sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape. In a particularly preferred embodiment, the plant is a maize plant.

The invention also provides microorganisms comprising a heterologous DNA sequence which encodes a protein of the VIP3 class, but preferably a VIP3A(c) protein. In a preferred embodiment, the microorganism is selected from the group consisting of bacteria, baculovirus, algae and fungi. In another preferred embodiment, the microorganism is selected from the group consisting of *Bacillus*, *Pseudomonas*, *Clavibacter*, and *Rhizobium*. Further encompassed by the invention are entomocidal compositions comprising microorganisms comprising a heterologous DNA sequence which encodes a protein of the VIP3 class, but preferably a VIP3A(c) protein.

The invention further relates to plants and microorganisms further comprising a second DNA sequence which encodes a second insecticidal protein. Particularly preferred second DNA sequences are those which encode a  $\delta$ -endotoxin, those which encode another protein of the VIP3 class, or those which encode a protein of the VIP1 or VIP2 classes. In a more preferred embodiment, the  $\delta$ -endotoxin is active against an insect selected from the group consisting of Lepidoptera and Coleoptera. In a more particularly preferred embodiment the  $\delta$ -endotoxin is active against *Ostrinia*, or *Diabrotica*. In another particularly preferred is a second DNA sequence which encodes a  $\delta$ -endotoxin protein selected from the group consisting of Cry1, Cry3, Cry5 and Cry9. In a more particularly preferred embodiment, the  $\delta$ -endotoxin is selected from the group consisting of Cry1Aa, Cry1Ab, Cry1Ac, Cry1B, Cry1C, Cry1D, Cry1Ea, Cry1Fa, Cry3A, Cry9A, Cry9C and Cry9B. Most particularly preferred are  $\delta$ -endotoxins selected from the group consisting of Cry1Ab, Cry1Ba and Cry9C proteins.

The invention further provides a method of controlling insects by contacting the insects with an insecticidal amount of a protein of the VIP3 class, but preferably a VIP3A(c) protein, or an insecticidal amount of a chemical ligand to a receptor of the VIP3 class of proteins. In

one preferred embodiment, the insects are contacted with a transgenic plant comprising a DNA sequence which expresses a protein of the VIP3 class, but preferably a VIP3A(c) protein. In another preferred embodiment, the insects are contacted with a an entomocidal composition comprising a protein of the VIP3 class, but preferably a VIP3A(c) protein, or comprising a DNA sequence which expresses a protein of the VIP3 class, but preferably a VIP3A(c) protein. In another preferred embodiment, the transgenic plant comprises a DNA sequence which expresses the VIP3A(a) protein. In another preferred embodiment the insect is selected from the group consisting of Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, and Acari. In a particularly preferred embodiment, the insect is a Coleoptera or Lepidoptera. In another particularly preferred embodiment, the insect is selected from the group consisting of black cutworm (*Agrotis ipsilon*), fall armyworm (*Spodoptera frugiperda*), beet armyworm (*S. exigua*), yellow striped armyworm (*S. ornithogalli*), southwestern corn borer (*Diatraea grandiosella*), sugarcane borer (*D. saccharalis*), corn earworm (*Helicoverpa zea*), mediterranean corn borer (*Sesamia nonagroides*), cabbage looper (*Trichoplusia ni*), velvetbean caterpillar (*Anticarsia gemmatilis*), diamondback moth (*Plutella xylostella*) and tobacco budworm (*Heliothis virescens*).

Also provided by the invention is a method of controlling insects wherein the transgenic plant or microorganism further comprises a second DNA sequence which encodes a second insecticidal protein such as those mentioned hereinbefore.

The invention further provides recombinant DNA sequences which encode a VIP3A(c) protein including homologues thereof. In another preferred embodiment, the DNA sequence is a synthetic sequence which has been altered for optimum expression in a plant, particularly where the DNA sequence has been optimized for expression in a maize plant. Also preferred are DNA sequences which comprise both a synthetic portion and a native portion. In a particularly preferred embodiment, the DNA sequence encoding the VIP3A(c) protein has been optimized for expression in a maize plant. Another preferred embodiment are DNA sequences which are homologous to a DNA sequence which encodes a VIP3A(c) protein. Particularly preferred are DNA sequences which hybridize under moderately stringent conditions to the *vip3A(c)* coding sequence. Yet another embodiment of the invention is a recombinant DNA sequence which expresses a protein of the VIP3 class, preferably a VIP3A(c) protein, under the control of a heterologous promoter,

or wherein the coding regions is incorporated into the genome of an organism where it is not naturally expressed or is expressed at higher levels than that occurring naturally.

The invention is further drawn to a method of identifying and isolating homologues of a VIP3A(c) protein, or of a DNA sequence which encodes said protein.

Also provided by the invention are expression cassettes comprising a promoter operably linked to a DNA sequence encoding a protein of the VIP3 class, but preferably a VIP3A(c) protein. In one preferred embodiment the promoter is selected from the group consisting of constitutive, tissue-preferred and tissue-specific promoters for expression in plants. In a particularly preferred embodiment, the promoter is selected from the group consisting of the ubiquitin, PEP carboxylase, LPT and MTL promoters. In another preferred embodiment, the promoter is functional in a microorganism.

The invention further provides a receptor to a protein of the VIP3 class and DNA sequences which. In one embodiment of the invention, the receptor comprises a death domain and a repeated EGF-motif. A more preferred embodiment of the invention comprises a receptor to the VIP3A(a). A more particularly preferred embodiment is the receptor protein sequence set forth in SEQ ID NO:9, and homologues thereto. Also encompassed by the invention are DNA sequences which encode these receptor proteins, e.g., the DNA sequence set forth in SEQ ID NO:8 and homologues thereto. The cDNA for the VIP3 receptor is contained in plasmid pCIB7113, which was deposited under the Budapest Treaty with the NRRL [Agricultural Research Service, Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 North University Street, Peoria, Illinois 61604, USA] on March 29, 1997 and has accession number B-21676. Antibodies to a receptor of the VIP3 class of proteins are also encompassed by the invention.

Also provided by the invention is a method of identifying a compound as a VIP3 receptor chemical ligand having pesticidal activity comprising exposing a cell, preferably an insect cell, to a test compound, and assaying said cell for apoptotic activity. In another embodiment of the invention, the method comprises measuring specific binding between VIP3 receptor and a test compound. A preferred embodiment are VIP3 receptor ligands identified by the method.

### ***Definitions***

"Plant pest" means any organism known to associate with plants and which, as a result of that association, causes a detrimental effect on the plant's health and vigor. Plant pests include but are not limited to fungi, bacteria, insects, and nematodes. The term plant

as used herein encompasses whole plants and parts of plants such as roots, stems, leaves and seed, as well as cells and tissues within the plants or plant parts.

The "VIP3 class of proteins" comprises VIP3A(a), VIP3A(b) VIP3A(c) and their homologues. "Homologue" is used throughout to mean that the indicated protein or polypeptide bears a defined relationship to other members of the VIP3 class of proteins. This defined relationship includes but is not limited to, 1) proteins which are at least 70%, more preferably 80% and most preferably 90% identical at the sequence level to another member of the VIP3 class of proteins while also retaining pesticidal activity, 2) proteins which are cross-reactive to antibodies which immunologically recognize another member of the VIP3 class of proteins, 3) proteins which are cross-reactive with a receptor to another member of the VIP3 class of proteins and retain the ability to induce programmed cell death, and 4) proteins which are at least 70%, more preferably 80% and most preferably 90% identical at the sequence level to the toxic core region of another member of the VIP3 class of proteins while also retaining pesticidal activity.

A "hybrid toxin" is used to indicate a genetic fusion, having domains operably linked so that, when translated, a functional chimeric protein is formed having, in the aggregate, the properties of the individual domains. "Domain" is used to indicate a region or portion of a protein or confers a recognizable function or structure which contributes to the overall functionality of the protein. It is recognized that a DNA sequence which encodes a protein domain is also encompassed by this definition.

"Heterologous" is used to indicate that a protein, polypeptide or nucleotide sequence has a different natural origin with respect to its current host. For example, if a *vip3A(a)* gene from a *Bacillus thuringiensis* is genetically transformed into a plant cell, then the gene is described as being heterologous with respect to its current host, which is the plant cell. Furthermore, if a *vip3A(a)* gene from *Bacillus thuringiensis* is genetically transformed into a *Pseudomonas* bacterium, then the gene is also described as being heterologous with respect to the *Pseudomonas*. "Heterologous" is also used to indicate that one or more of the domains present in a chimeric protein, polypeptide or nucleotide sequence differ in their natural origin with respect to other domains present. For example, if the toxic domain from VIP3A(a) protein is fused to the binding domain from the VIP1A(a) protein to make a functional insecticidal protein, then the chimeric fusion would have domains that are heterologous to each other. In addition, a heterologous chimeric protein or polypeptide comprising the fusion of a toxic domain from VIP3A(a) protein to the binding domain from

the VIP1A(a) protein, when expressed in a plant, would also be considered heterologous with respect to the plant host.

The term "chimeric" is used to indicate that the protein, polypeptide, or nucleotide sequence is comprised of domains at least one of which has an origin that is heterologous with respect to the other domains present. These chimeric proteins or polypeptides are encoded by chimeric nucleotide sequences which have been fused or ligated together resulting in a coding sequence which does not occur naturally. Such chimeric constructions may also be designated as "recombinant."

"Expression cassette" as used herein means a DNA sequence capable of directing expression of a gene in plant cells, comprising a promoter operably linked to an amino acid coding region which is operably linked to a termination region. The gene may be chimeric, meaning that at least one component of the gene is heterologous with respect to at least one other component of the gene. The gene may also be naturally occurring, but which has been obtained in a recombinant form useful for genetic transformation of a plant or microorganism.

### **Figures**

**Figure 1:** Amino acid sequence of the receptor for VIP3A(a) translated from the cDNA. Several features of the protein are shown: *dotted line* - signal peptide (amino acid 13 to 35); *italic* - domain spanning the putative death domain (amino acid 81-205); *double underline* - sequences with strong homology to sequences found in consensus death domains; *bold* - CKC motif repeated six times spanning the EGF-motifs; *underline* - sequences repeated within the EGF-motifs.

### **Arthropod Pests**

For purposes of the present invention, pests include insects and arachnids selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, and Acari, particularly Coleoptera and Lepidoptera.

A list of pests associated with major crop plants are provided, for example, in Tables 1-10 on pages 13 to 20 of WO 96/10083, which is incorporated herein by reference. Further pests are provided in the following Tables 1 - 8. Such pests are included within the scope of the present invention.

Table 1: Lepidoptera (Butterflies and Moths)

Maize <i>Sesamia nonagroides</i> , Mediterranean corn borer <i>Ostrinia furnacalis</i> , Asian corn borer	Crucifers (broccoli, cabbage, cauliflower, collards) <i>Artogeia rapae</i> , imported cabbageworm  <i>Pieris brassicae</i> , cabbage butterfly <i>Trichoplusia ni</i> , cabbage looper <i>Plutella xylostella</i> , diamondback moth <i>Spodoptera exigua</i> , beet armyworm <i>Agrotis ipsilon</i> , black cutworm <i>Agrotis segetum</i> , common cutworm <i>Mamestra configurata</i> , bertha army worm
Cotton <i>Helicoverpa armigera</i> , cotton bollworm	
Rice <i>Chilo suppressalis</i> , asiatic rice borer <i>Scirpophaga</i> sp.	
Tomato  <i>Helicoverpa zea</i> , tomato fruitworm <i>Spodoptera exigua</i> , beet armyworm <i>Spodoptera frugiperda</i> , fall armyworm <i>Spodoptera ornithogalli</i> , yellowstriped armyworm <i>Spodoptera praefica</i> , western yellowstriped armyworm <i>Spodoptera eridania</i> , southern armyworm <i>Agrotis ipsilon</i> , black cutworm <i>Peridroma saucia</i> , variegated cutworm <i>Papaipema nebris</i> , stalk borer <i>Trichoplusia ni</i> , cabbage looper <i>Keiferia lycopersicella</i> , tomato pinworm <i>Manduca sexta</i> , tobacco hornworm <i>Manduca quinquemaculata</i> , tomato hornworm	Grapes <i>Endopiza viteana</i> , grape berry moth  Deciduous Fruits and Nuts <i>Cydia pomonella</i> , codling moth <i>Platynota idaeusalis</i> , tufted apple bud moth  Peppers  <i>Ostrinia nubilalis</i> , European corn borer <i>Spodoptera exigua</i> , beet armyworm <i>Spodoptera eridania</i> , southern armyworm  Potato <i>Ostrinia nubilalis</i> , European corn borer <i>Phthorimaea operculella</i> , potato tuberworm  Canola <i>Plutella xylostella</i> , diamondback moth  Sugarcane <i>Diatraea saccharalis</i> , sugarcane borer

Table 2: Coleoptera (Beetles)

Rice <i>Oulema oryzae</i> , rice beetle	Potato <i>Leptinotarsa decemlineata</i> , Colorado potato beetle <i>Epitrix cucumeris</i> , potato flea beetle <i>Hemicrepidus memnonius</i> , wireworms <i>Melanotus</i> spp., wireworms
Tomato <i>Leptinotarsa decemlineata</i> , Colorado potato beetle <i>Epitrix hirtipennis</i> , tobacco flea beetle	

Crucifers (broccoli, cabbage, cauliflower,  
collards)

*Phyllotreta cruciferae*, crucifer flea beetle  
*Phyllotreta pusilla*, western black flea beetle

Canola

*Ceutorhynchus assimilis*, cabbage seedpod weevil

*Phyllotreta cruciferae*, crucifer flea beetle

Peppers

*Anthonomus eugenii*, pepper weevil

**Table 3: Homoptera (Whiteflies, Aphids etc..)**

Rice	Melon
<i>Nilaparvata lugens</i>	<i>Bemisia argentifolii</i> , silverleaf whitefly
<i>Sogatella furcifera</i>	<i>Bemisia tabaci</i> , sweetpotato whitefly
<i>Laodelphaax striatellus</i>	
Tomato	Carrot
<i>Myzus persicae</i> , green peach aphid	<i>Cavariella aegopodii</i> , carrot aphid
<i>Macrosiphum euphorbiae</i> , potato aphid	
<i>Trialeurodes vaporariorum</i> , greenhouse whitefly	Canola
<i>Bemisia tabaci</i> , sweetpotato whitefly	<i>Brevicoryne brassicae</i> , cabbage aphid
<i>Bemisia argentifolii</i> , silverleaf whitefly	
Crucifers (broccoli, cabbage, cauliflower, collards)	Vegetables
<i>Brevicoryne brassicae</i> , cabbage aphid	<i>Aphis fabae</i> , bean aphid
<i>Myzus persicae</i> , green peach aphid	
Peppers	Sugar Beet
<i>Myzus persicae</i> , green peach aphid	<i>Pemphigus popullivivae</i> , sugar beet root aphid
Potato	Deciduous Fruits and Nuts
<i>Empoasca fabae</i> , potato leafhopper	<i>Dysaphis plantaginea</i> , rosy apple aphid
<i>Myzus persicae</i> , green peach aphid	Sugarcane
<i>Macrosiphum euphorbiae</i> , potato aphid	<i>Saccharosydne saccharivora</i> , West Indian cane fly
<i>Paratrioza cockerelli</i> , potato psyllid	<i>Sipha flava</i> , yellow sugarcane aphid

**Table 4: Hemiptera (Bugs)**

Tomato
lygus bug
<i>Acrosternum hilare</i> , green stink bug
<i>Euschistus servus</i> , brown stick bug

**Table 5: Orthoptera (Grasshoppers, Crickets, and Cockroaches)**

Wheat
<i>Melanoplus sanguinipes</i> , migratory grasshopper



**Table 6: Diptera (Flies and Mosquitoes)**

Tomato	Carrot
<i>Liriomyza trifolii</i> , leafminer	<i>Psilia rosae</i> , carrot rust fly
<i>Liriomyza sativae</i> , vegetable leafminer	Sugarbeet
<i>Scrobipalpula absoluta</i> , tomato leafminer	<i>Tetanops myopaeformis</i> , sugarbeet root maggot
Crucifers (broccoli, cabbage, cauliflower, collards)	Vegetables
<i>Delia brassicae</i> , cabbage maggot	<i>Liriomyza sativae</i> , vegetable leaf miner
<i>Delia radicum</i> , cabbage root fly	

**Table 7: Thysanoptera (Thrips)**

Tomato	Peppers
<i>Frankliniella occidentalis</i> , western flower thrips	<i>Thrips palmi</i> , melon thrips
<i>Frankliniella fusca</i> , tobacco thrips	
<i>Thrips tabaci</i> , onion thrips	Potato
	<i>Thrips palmi</i> , melon thrips
Crucifers (broccoli, cabbage, cauliflower, collards)	
<i>Thrips tabaci</i> , onion thrips	

**Table 8: Acari (Mites and Ticks)**

Tomato
<i>Tetranychus urticae</i> , two-spotted spider mite
<i>Aculops lycopersici</i> , tomato russet mite
<i>Steneotarsonemus pallidus</i> , cyclamen mite
Citrus
<i>Panonychus citri</i> , citrus red mite
<i>Brevipalpus lewisi</i> , citrus flat mite
<i>Phyllocoptruta oleivora</i> , citrus rust mite
Deciduous Fruits and Nuts
<i>Panonychus ulmi</i> , European red mite
<i>Tetranychus sp.</i> , spider mite

For purposes of the present invention, pests also include fungal phytopathogens of plants. A list of fungal pests associated with major crop plants is provided in Table 9. Such pests are included within the scope of the present invention.

**Table 9: Fungal Diseases of Plants****Ear Molds**

Gibberella ear mold	<i>Gibberella zeae</i> <i>G. saubineti</i>
Aspergillus ear rot	<i>Aspergillus flavus</i> <i>A. parasiticus</i>
Diplodia ear rot	<i>Diplodia maydis</i> <i>D. macrospora</i>
Fusarium ear rot	<i>Fusarium moniliforme</i> <i>F. monilif. var. subglutinans</i>

**Stalk Rots**

Pythium stalk rot	<i>Pythium aphanidermata</i>
Anthracnose stalk rot	<i>Colletotrichum graminicola</i> <i>C. tucumanensis</i> <i>Glomerella graminicola</i>
Diplodia stalk rot	<i>Diplodia maydis</i> <i>D. zeae-maydis</i> <i>Stenocarpella maydis</i> <i>Macrodiplodia zeae</i> <i>Sphaeria maydis</i> <i>S. zeae</i> <i>D. macrospora</i>
Fusarium stalk rot	<i>Fusarium moniliforme</i>
Gibberella stalk rot	<i>G. zeae</i> <i>G. saubineti</i>

**Stewart's wilt & leaf blight***Erwinia stewartii***Leaf Diseases**

Northern corn leaf blight	<i>Exserohilum turcicum</i>
Southern corn leaf blight	<i>Bipolaris maydis</i>
Gray leaf spot	<i>Cercospora zeae-maydis</i> <i>C. sorghi var. maydis</i>
Anthracnose leaf blight	<i>Colletotrichum graminicola</i>
Common rust	<i>Puccinia sorghi</i> <i>P. maydis</i>
Southern rust	<i>Puccinia polysora</i> <i>Dicaeoma polysorum</i>
Head smut	<i>Sphacelotheca reiliana</i>
Common smut	<i>Ustilago maydis</i>
Carbonum leaf spot	<i>Helminthosporium carbonum</i>
Eye spot	<i>Kabatiella zeae</i>

**Downy Mildews**

Sorghum downy mildew	<i>Peronosclerospora sorghi</i>
----------------------	---------------------------------

Brown stripe downy mildew	<i>Sclerophthora rayssiae</i>
Sugarcane downy mildew	<i>Peronosclerospora sacchari</i>
Phillipine downy mildew	<i>Peronoscler. philippinensis</i>
Java downy mildew	<i>Peronosclerospora maydis</i>
Spontaneum downy mildew	<i>Peronosclerospora spontanea</i>
Rajasthan downy mildew	<i>Peronosclerospora heteropogoni</i>
Graminicola downy mildew	<i>Sclerospora graminicola</i>
Rusts	<i>Puccinia graminis f.sp. tritici</i> <i>Puccinia recondita f.sp. tritici</i> <i>Puccinia striiformis</i>
Smuts	<i>Tilletia tritici</i> <i>Tilletia controversa</i> <i>Tilletia indica</i> <i>Ustilago tritici</i> <i>Urocystis tritici</i>
Root rots, Foot rots and Blights	<i>Gaeumannomyces graminis</i> <i>Pythium spp.</i> <i>Fusarium culmorum</i> <i>Fusarium graminearum</i> <i>Fusarium avenaceum</i> <i>Drechslera tritici-repentis</i> <i>Rhizoctonia spp.</i> <i>Colletotrichum graminicola</i> <i>Helminthosporium spp.</i> <i>Microdochium nivale</i> <i>Pseudocercospora herpotrichoides</i>
Mildews	<i>Erysiphe graminis f.sp. tritici</i> <i>Sclerophthora macrospora</i>
Miscellaneous Fungal Diseases	<i>Septoria tritici</i> <i>Septoria nodorum</i>

The proteins of the VIP3 class are secreted to the media by *Bacillus spp.* in vegetative stages of growth. VIP3A(a) is a member of a newly discovered class of proteins displaying insecticidal activity against a broad spectrum of lepidopteran insects including black cutworm (*Agrotis ipsilon*), fall armyworm (*Spodoptera frugiperda*), beet armyworm (*S. exigua*), yellow striped armyworm (*S. ornithogalli*), southwestern corn borer (*Diatraea grandiosella*), sugarcane borer (*D. saccharalis*), corn earworm (*Helicoverpa zea*), Mediterranean corn borer (*Sesamia nonagroides*), cabbage looper (*Trichoplusia ni*), velvetbean caterpillar (*Anticarsia gemmatilis*), diamondback moth (*Plutella xylostella*) and

tobacco budworm (*Heliothis virescens*). Some of these lepidopteran insects have been shown to be very resistant to other insecticidal proteins such as  $\delta$ -endotoxin. For example, the reported LC<sub>50</sub> for Cry1A(c), which is one of the most effective  $\delta$ -endotoxin against black cutworm, is greater than 6000 ng/cm<sup>2</sup> (MacIntosh *et al.*, J. Invertebr. Pathol. 56:258-266 (1990)). In contrast, it takes 260-fold less of VIP3A(a) protein to kill 50% of the black cutworm larvae. Thus, the VIP3A(a) protein displays a unique spectrum of insecticidal activities.

The present invention provides a new member of the VIP3 class of proteins, the VIP3A(c) protein isolated from strain AB51 (pCIB7112 deposited on March 28, 1997 as Accession No. NRRL B-21675; all deposits were made in accordance with the Budapest Treaty by submission to the Agricultural Research Service, Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 North University Street, Peoria, Illinois 61604, USA) as disclosed in SEQ ID NO:5-6..

The VIP3A(c) protein encoding DNA sequence was identified and isolated through the use of PCR technology. In particular, primer sequences can be made which recognize either conserved or variable regions of the coding sequence, and are then used to screen DNA samples obtained from either known or unknown strains.

It is recognized that there are multiple approaches to identifying and isolating homologues within the VIP3 class of proteins and the DNA sequences which encode them, which approaches are well known to those skilled in the art.

The DNA and protein sequences for the VIP3A(a) and VIP3A(c) proteins are aligned in Table 12.

**Table 12: Alignment of VIP3A(a) (Upper Line) against VIP3A(c) (Lower Line)**

```

1 MNKNNTKLSTRALPSFIDYFNIGIYGFATGIKDIMNMIFKTDGTGGDLTLDE 50 SEQ ID NO:2
|||||.|||||||
1 MNKNNAKLSTRALPSFIDYFNIGIYGFATGIKDIMNMIFKTDGTGGDLALDE 50 SEQ ID NO:6
51 ILKNQQLLNDISGKLDGVNGSLNDLIAQGNLNTELSKEILKIANEQNQVL 100
|||.|||||||
51 ILENQQLLNDISGKLDGVNGSLNDLIAQGNLNTELSKEILKIANEQNQVL 100
101 NDVNNKLD AINTMLRVYLPKITSMLS DVMKQNYALSLQIEYLSKQLQEIS 150
|||||
101 NDVNNKLD AINTMLRVYLPKITSMLS DVMKQNYALSLQIEYLSKQLQEIS 150
151 DKLDIINVNLINSTL TEITPAYQRIKYVNEKFEELTFATETSSKVKKDG 200

```

```

151 DKLDIINVNLINSLTEITPAYQRIKYVNEKFEELTFATETSSKVKKDG 200
201 SPADILDELTELTELAKSVTKNDVDGFEFYLNTHDVMVGNNLFGRSALK 250
201 SPADIRDELSELTELAKSVTQNDVDGFEFYLNTHDVMVGNNLFGRSALK 250
251 TASELITKENVKTSSEVGNVNFILVLTALQAQAFLLTTCRKLLGLAD 300
251 TASELITKENVKTSSEVGNVNFILVLTALQAQAFLLTPCRKLLGLAD 300
301 IDYTSIMNEHLNKEKEEFRVNILPTLSNTFSNPYAKVKGSEDEAKMIVE 350
301 IDYTSIMNEHLNKEKEEFRVNILPTLSNTFSNPYAKVKGSEDEAKMIVE 350
351 AKPGHALIGFEISNDSITVLKVYEAKLKQNYQVDKDSLSEVIYGDMDKLL 400
351 AKPGHALIGFEISNDSITVLKVYEAKLKQNYQVDKDSLSEVIYGDMDKLL 400
401 CPDQSEQIYYTNNIVFPNEYVITKIDFTKKMKTLYEVTANFYDSSTGEI 450
401 CPDQSGQIYYTNNIVFPNEYVITKIDFTKKMKTLYEVTANFYDSSTGEI 450
451 DLNKKKVESSEAEYRTLSANDDGVYMLPGVISEFTLTPINGFGLQADENS 500
451 DLNKKKVESSEAEYRTLSANDDGVYMLPGVISEFTLTPINGFGLQADENS 500
501 RLITLTCKSYLRELLLATDLNKETKLIVPPSGFISNIVENGSIEDNLE 550
501 RLITLTCKSYLRELLLATDLNKETKLIVPPSGFISNIVENGSIEDNLE 550
551 PWKANNKNAYVDHTGGVNGTKALYVHKDGGISQFIGDKLKPKTEYVIQYT 600
551 PWKANNKNAYVDHTGGVNGTKALYVHKDGGISQFIGDKLKPKTEYVIQYT 600
601 VKGKPSIHLKDENTGYIHYEDTNNNLEDYQTINKRFTTGTDLKGVYLILK 650
601 VKGKPSIHLKDENTGYIHYEDTNNNLEDYQTINKRFTTGTDLKGVYLILK 650
651 SQNGDEAWGDNFIILEISPSEKLLSPELINTNNWTSTGSTNISGNTLTLY 700
651 SQNGDEAWGDNFIILEISPSEKLLSPELINTNNWTSTGSTNISGNTLTLY 700
701 QGGRGILKQNLQLDSFSTYRVYFSVSGDANVRIRNSREVLFEKRYM 746
701 QGGRGILKQNLQLDSFSTYRVYFSVSGDANVRIRNSREVLFEKKDI 746

```

### ***Polypeptide Domains of the VIP3 Class of Proteins***

It has been shown that the VIP3A(a) protein undergoes proteolytic processing when mixed with the gut fluids of insect larvae. When gut fluids isolated from black cutworm are mixed with purified VIP3A(a), four major proteolytic products derived from VIP3A(a) can be identified having a molecular weight of approximately 66, 45, 33 and 22 kDa. The 22 kDa band comprises the N-terminal portion of the VIP3A(a) protein from amino acid 1 to amino acid 198 of SEQ ID NO:2. The 66 kDa band comprises the rest of the VIP3A(a) protein from

amino acid 200 to amino acid 789 of SEQ ID NO:2. Both the 45 and 33 kDa bands are derived by proteolysis from the 66 kDa band and constitute amino acid 412 to amino acid 789, and from amino acid 200 to amino acid 455, respectively, of SEQ ID NO:2. The 33 kDa band is the main component of the VIP3A(a) protein that remains after an incubation period of more than two hours. This 33 kDa "toxic core" domain (amino acids 200 to 455 of SEQ ID NO:2) of the VIP3A(a) protein retains full insecticidal properties against a broad spectrum of lepidopteran insects. Similar results are obtained when VIP3A(a) is incubated with gut fluids isolated from fall armyworm, another insect sensitive to VIP3A(a).

In addition to the toxic core domain, the VIP3A(a) protein also possesses a stabilizing domain at the C-terminus. The role of the stabilizing domain was explored using mutants of the VIP3A(a) protein and the VIP3A(c) protein, neither of which display insecticidal properties when ingested by insects known to be sensitive to VIP3A(a). When similar studies addressing the stability in black cutworm gut fluid was conducted with VIP3A(a)-mutants, in particular with a mutant of the VIP3A(a) protein that contains three point mutations located at the carboxy-terminal domain (amino acid 742 (E→D); amino acid 770 (S→P); and amino acid 784 (Y→H)), it was found that the protein was completely hydrolyzed. Similar results were obtained for the VIP3A(c) (SEQ ID NO:6) protein isolated from AB51, which shares an overall identity of 96% with the VIP3A(a) protein but lacks the carboxy-terminal domain of VIP3A(a). Both the mutant and VIP3A(c) protein, however, are active against the insect cell line Sf-9. These results indicate that the function of the carboxy-terminal domain of proteins of the VIP3 class is to provide stability to the protein in the gut environment of susceptible insects.

#### ***Hybrid Toxins Comprising a VIP3 Region and a Heterologous Region***

Toxins, enzymes, transcription factors, antibodies, cell binding moieties or other protein domains can be operably linked to the novel proteins of the present invention by producing in frame genetic fusions which, when translated by ribosomes, would produce a fusion protein with the combined attributes of the VIP and the other component used in the fusion. Furthermore, if the protein domain fused to the VIP has an affinity for another protein, nucleic acid, carbohydrate, lipid, or other chemical or factor, then a three-component complex can be formed. This complex will have the attributes of all of its components. A similar rationale can be used for producing four or more component complexes. These complexes are useful as insecticidal toxins, pharmaceuticals, laboratory

reagents, and diagnostic reagents, etc. Examples where such complexes are currently used are fusion toxins for potential cancer therapies, reagents in ELISA assays and immunoblot analysis.

The hybrid toxins of the invention include chimeric proteins having a toxic core domain which is heterologous to the stabilizing domain. Hybrid toxins are also created by combining an antibody, or immunologically-active fragment thereof, which immunologically recognizes the VIP3 receptor with a toxic domain from other proteins. The toxin domain is obtained from a number of cytotoxic proteins. These include but are not limited to *Bacillus* toxins, including endotoxins and vegetative insecticidal proteins. See for example US application serial no. 08/037,057, filed March 25, 1993 and U.S. application serial no. 07/951,715 filed September 25, 1992, herein incorporated by reference. Other toxins include catalytic ribosome inactivators such as gelonin, *Pseudomonas* exotoxin A or phytoalexin, (the structure of *Pseudomonas* exotoxin has been well characterized in Chaudhary *et al.*, J. Biol. Chem. 265:16303-16310 (1990)); cell metabolism disrupters, such as ribonucleases, (see, for example, Mariani *et al.* Nature 347:737-741 (1990)); Barnase toxin (or PE-Bar), a chimeric toxin derived from *Pseudomonas* exotoxin A and a ribonuclease, (see, Prior *et al.* Cell 64:1017-1023 (1991)); hydrophilic peptides that create pores in membranes (see, Frohlich and Wells, Int. J. Peptide Protein Res. 37:2-6 (1991)).

#### **Mode of Action of VIP3A(a)**

The VIP3A(a) protein has been shown to be active against a broad spectrum of plant pests. For example, histopathological observations indicate that VIP3A(a) ingestion by susceptible insects such as black cutworm (*Agrotis ipsilon*) and fall armyworm (*Spodoptera frugiperda*) causes gut paralysis at concentrations as low as 4 ng/cm<sup>2</sup> of diet, with complete lysis of the gut epithelial cells resulting in larval death at concentrations above 40 ng/cm<sup>2</sup>. Less susceptible insects like European corn borer (*Ostrinia nubilalis*) do not develop any pathology upon ingesting VIP3A(a). While the proteolytic processing of the VIP3A(a) protein by midgut fluids obtained from susceptible and non-susceptible insects is comparable, *in vivo* immuno-localization studies show that VIP3A(a) binding is restricted to gut cells of susceptible insects. Therefore, the insect host range for VIP3A(a) seems to be determined by its binding ability to gut cells. Histopathological observations indicate that midgut epithelial cells of susceptible insects are the primary target for the VIP3A(a) insecticidal protein and their subsequent lysis is the primary mechanism of lethality.

Programmed cell death is an active process of self-destruction that seems to be important for development and maintenance of multicellular organisms (Clem, R.J. *et al.* Science 254: 1388-1390 (1991)). Cells undergoing apoptosis, which is a form of programmed cell death, generate membrane-bound apoptotic bodies and activate endogenous nucleases that cleaves the chromatin into discrete fragments. SF-9 insect cells derived from *S. frugiperda* exposed to the VIP3A(a) protein undergo a series of cytological and molecular changes including membrane protrusions, profuse vacuolization and endonucleolysis which are indicative of an apoptotic-type of programmed cell death. Histological studies have shown that the VIP3A(a) protein targets midgut epithelial cells of susceptible insects initiating a series of cytological changes comprising profuse vacuolization and swelling prior to cell lysis and larval death. These midgut cells also experienced an endonucleolysis process when exposed to the VIP3A(a) protein as revealed by *in situ* detection of DNA fragmentation. These results indicate that VIP3A(a) exerts its insecticidal properties on susceptible insect cells by triggering an apoptotic-type of programmed cell death.

#### ***The Receptor for VIP3A(a) has been Isolated***

The immunohistochemistry results provided above indicate that VIP3A(a) has the ability to bind to the apical membranes of midgut epithelial cells and that this binding triggers the process that will eventually end with cell lysis. This indicates that there exists one or more proteins located in the apical membrane that recognize and bind to VIP3A(a) acting as a receptor. This receptor signals the interaction with VIP3A(a) and triggers the process of apoptosis. Thus, the receptor will mediate the response of the insect cell to VIP3A(a).

To isolate this receptor, a cDNA library was screened which was made from mRNA isolated from midgut tissue of black cutworm. The objective of the screen was to identify and isolate cDNA sequences which encode proteins that will interact with VIP3A(a) in the two hybrid system (see Fields, S. and Song, O.-K. Nature 340:245-246 (1989)). This approach resulted in the identification and isolation of one cDNA whose encoded protein strongly interacted with the VIP3A(a) protein. This 1.75 Kb-long cDNA (SEQ ID NO:8) encodes a protein of approximately 48 kDa (396 amino acids; see SEQ ID NO:9). The cloned cDNA is similar in size to the mRNA encoding the cDNA as analyzed by Northern. A portion of the DNA sequence which encodes the first 5 to 20 amino acids may be missing. The following features can be identified in the cDNA encoded protein (see Figure 1): 1) it



contains a signal peptide; 2) it contains a domain with homology to the so-called death domain (Feinstein, E. *et al.* Trends in Biochem. 20:342-344 (1995)); and 3) it contains EGF-like motifs or repeats (Fantl, W.J. *et al.* Annu. Rev. Biochem. 62:453-481 (1993)). A search of protein databases using the receptor of VIP3A(a) showed homology with a family of extracellular glycoproteins known as Tenascins (Pearson, C.A. *et al.* EMBO J. 7:2677-2681 (1988)) or Hexabrachion (Nies, D.E. *et al.* J. Biol. Chem. 266:2818-2823 (1991)). This family of proteins contains EGF-like repeats, interacts with multiple ligands, and performs a role in cell adhesion and/or signaling. The combination of a death domain and repeated EGF-motifs as observed in the VIP3 receptor is unique among programmed cell death receptors.

In addition, a portion of the VIP3A(a) receptor shares homology with the so-called "death domain." The death domain is a 60 to 70 amino acid long motif which is involved in protein to protein interaction and is shared by proteins with diverse cellular functions (Feinstein, E. *et al.* Trends in Biochem. 20:342-344 (1995)). Some of the protein members containing death domain motifs include receptors known to be associated with apoptotic processes. Some examples include the Fas receptor (Brakebush, C. *et al.* EMBO J. 11:943-950 (1992)) and the tumor necrosis factor (TNF) (Tartaglia, L.A. *et al.* Cell 74:845-853 (1993)).

Homologues to the VIP3A(a) receptor can be identified and isolated by various means, for example, by nucleic acid hybridization. Southern blot analysis can be performed on DNA samples taken from insect cells or fungal cells that has been enzyme restricted, run in agarose and blotted onto nitrocellulose and/or nylon filters. The Southern blot can be probed with the full-or partial length of the nucleic acid encoding the receptor of the VIP3A(a) protein under low stringency hybridization and washing conditions. The genes can be readily cloned and sequenced from a cDNA or genomic library. A size-selected genomic library can also be obtained to facilitate cloning of the gene(s) of interests. The technical protocols to perform the experiments outlined above are readily available (see for instance Molecular Cloning, A Laboratory Manual, Second Edition, Vols. 1-3, Sambrook *et al.* (eds.) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) and reference therein).

#### ***Antibodies to VIP3A(a) and Its Receptor***

Polyclonal and monoclonal antibodies to a VIP3 protein or its receptor, including fragments thereof which immunologically recognize a portion of either protein, are provided. The antibody and monoclonal antibodies of the present invention can be prepared by utilizing a VIP3 protein or its receptor as the antigen.

The antibodies of the invention include polyclonal and monoclonal antibodies as well as fragments thereof which retain their ability to bind a VIP3 protein or its receptor. An antibody, monoclonal antibody, or fragment thereof is said to be capable of binding a molecule if it is capable of specifically reacting with the molecule to thereby bind the molecule to the antibody, monoclonal antibody, or fragment thereof. The term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to include intact molecules as well as fragments or binding regions or domains thereof (such as, for example, Fab and F(ab)<sub>2</sub> fragments) which are capable of binding hapten. Such fragments are typically produced by proteolytic cleavage, such as papain or pepsin. Alternatively, hapten-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

Methods for the preparation of the antibodies of the present invention are generally known in the art. For example, see *Antibodies, A Laboratory Manual*, Ed Harlow and David Lane (eds.) Cold Spring Harbor Laboratory, NY (1988), as well as the references cited therein. Standard reference works setting forth the general principles of immunology include: Klein, J. *Immunology: The Science of Cell-Noncell Discrimination*, John Wiley & Sons, NY (1982); Dennett, R., *et al.* *Monoclonal Antibodies, Mybridoma: A New Dimension in Biological Analyses*, Plenum Press, NY (1980); and Campbell, A. "Monoclonal Antibody Technology," In *Laboratory Techniques in Biochemistry and Molecular Biology*, Vol. 13, Burdon *et al.* (eds.), Elsevier, Amsterdam (1984). See also, US Patent Nos: 4,609,893; 4,713,325; 4,714,681; 4,716,111; 4,716,117; and 4,720,459.

It is recognized that following the methods described herein, antibodies specific for a particular VIP3 protein or its receptor can be generated. The subset of MAb lines which possess the desired binding specificity can be used as a source of messenger RNA for cloning of the cDNA for the particular monoclonal antibody.

The cloned DNA can then be sequenced by methods known in the art. See, for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. Edition, Cold Spring Harbor Laboratory Press, NY (1989) vol. 1-3, and the references cited therein. From the nucleic acid sequence, the protein sequence of the binding region from the selected MAb can be deduced.

One use of the antibodies and monoclonal antibodies of the invention includes but is not limited to the production of hybrid toxin molecules. That is, when linked, the monoclonal antibody or antibody fragment retains its binding properties and the toxin moiety retains its cytotoxic properties.

Various methods are known for obtaining antibody genes. One method is to clone a random library of antibody genes in a phage and screen the library for ability to bind to a VIP3 protein or its receptor. Another available approach is to generate monoclonal antibodies which bind to a VIP3 protein or its receptor and then clone the antibody genes from such lines. For the present example, the second method is used. Antibody genes can be cloned from hybridoma cells using primers to conserved DNA sequences within the constant regions and the framework regions of the variable regions and amplified for cloning using the polymerase chain reaction (PCR). See generally, Mullis *et al.*, *Meth. Enzymol.*, 155:335-350 (1987); Erlich, (ed.), *PCR Technology*, Stockton Press (New York 1989). A database of mouse heavy chain and light chain sequences compiled by Kabat *et al.*, US Dept Health and Human Services, US Government Printing Offices (1991) has been successfully used to generate both isotype specific and degenerate primers for cloning antibody genes. (Jones *et al.* *Bio/technology* 9:88-89 (1991)). Additionally, techniques are well known for cloning of smaller fragments of antibodies (Fab) which possess the binding properties of the original antibody. Complete antibodies are large molecules (150 kDa), but much smaller Fab and Fv antigen-binding fragments (12 kDa-50 kDa) have been shown to retain full binding affinity. Single chain Fv fragments (scFv) in which Vh and Vl domains are linked by a hydrophilic and flexible peptide have been used successfully to target enzymes and toxins to specific cells (Bird, *Science* 423:423-426 (1988); Huston, *PNAS* 85:5879-5883 (1988)). Single Vh domains (Dabs) and single complementary determining regions as small as 20 amino acids in length, called minimal recognition units (m.r.u.), have also been used for antigen binding (Ward, *Nature* 341:544-546 (1989); Taub, *J. Biol. Chem* 264:259-265 (1989); Williams, *PNAS* 86:5537-5541 (1989)). Thus, it is possible to reduce the binding domain specific for a VIP3 or its receptor to a very small size.

Polymerase chain reaction technology and specific oligonucleotide primers are used to clone immunoglobulin genes or regions from immunoglobulin genes. PCR primers specific for both the heavy and light chains of IgM and the three IgG isotypes were selected from the Kabat database described above. Primers for the region encoding the NH<sub>2</sub>-terminal end of the mature variable region were designed to initiate at the first framework region and were made with some degeneracy to allow these to be used as "universal primers". The 3'

primers used for the specific PCR amplification of the variable regions were designed from conserved sequences of the first constant domain (CH1) of both the light and heavy chains. A different 3' primer is used for immunoglobulin isotypes IgG1, IgG3, and IgM. Isotypes IgG2A and IgG2B can be amplified with the same primers used for IgG1. Antibody variable regions are cloned into a light and heavy chain expression vector containing an endoplasmic reticulum signal peptide and the constant regions of IgG1 light and heavy chains, respectively.

Primer sequences used for the PCR cloning of the mouse immunoglobulin light and heavy variable regions are available in the published literature (Coloma *et al.* Bio/Techniques 11: 152-156 (1991); Jones *et al.* Bio/Technology 9:88-89 (1991)). Oligonucleotides were made on an Applied Biosystems DNA synthesizer 380B (Applied Biosystems, Foster City, CA) using standard conditions as described below. The PCR primers incorporate restriction sites and, after amplification and digestion, can be cloned into a plant expression vector under the control of a plant-expressible promoter. Restriction sites were chosen that were known to be absent in sequenced antibody genes.

Another use of the polyclonal and/or monoclonal antibodies of the invention includes the stimulation of apoptosis by targeting the receptor to Vip3A with antibodies. The interaction of antibodies raised against cell surface-located proteins that are involved in controlling the cell growth result in the induction of apoptosis by means of preventing the said receptor from binding to its natural ligand(s). For instance, the anti-APO-1 antibody completely blocks proliferation of leukemia cells bearing the APO-1 protein and triggers apoptosis in these cells (Trauth, B.C. *et al.* Science 245:301-305 (1989)). Also, the activity resulting from the interaction between a given receptor and a ligand is mimicked by substituting the ligand for antibodies raised against the receptor. For instance, the addition of certain anti-Fas antibodies to cells bearing the Fas receptor in their cell surfaces will mediate apoptosis in a similar fashion as when the ligand of the Fas receptor is added (Itoh, N. *et al.* Cell 66:233-243 (1991)).

The receptor to Vip3A(a) isolated from black cutworm shares homology with a family of extracellular glycoproteins known as Tenascins, and in particular with Tenascin-X (Bristow, J. *et al.* J. Cell Biol. 122:265-278 (1993)). Tenascin-Xs are known to be involved in cell-to-cell adhesion and signaling. Lack of functionality of Tenascin-X either by mutation or by removal of the gene leads to lethality. Therefore, antibodies raised against different domains of the receptor to Vip3A(a) either effectively block the receptor from binding to its ligand(s) or mimic the interaction of the Vip3A(a) protein triggering apoptosis. This approach

is extended to different receptors with similar biological functions. In this sense, antibodies raised against insect cell receptors involved in crucial cell growth and interaction processes lead to induction of apoptosis and are used as a strategy to control insects.

***Screening for Novel Insecticidal Activities whose Mode of Action is Apoptosis***

The materials described in this invention are used to screen for chemical ligands that have pesticidal properties triggering apoptotic responses. Chemical ligands include small organic molecules, peptides, and proteins. In one embodiment of the invention, insect cell lines are used as model organisms for insects to screen for compounds that are insecticidal as a consequence of their ability to induce apoptosis. These cell lines are handled in a high-throughput screening format where the cells are grown in multi-well plates and are exposed to a variety of compounds. Yeast is also used as a model organism. Using procedures described herein or known in the art, determining whether a compound is pesticidal as a consequence of inducing apoptosis is accomplished.

One means by which to identify compounds that trigger apoptotic responses through interaction with a known receptor is to resort to identified receptors involved in the signal transduction pathway triggered in apoptotic insect cell lines. These receptors are transformed into heterologous cell lines creating isogenic lines with one of them containing a gene for expression of a specific receptor and another one which does not either possess, or express, such a gene. These cell lines are handled in a high-throughput screening format whereby the transformed cell lines expressing the receptor have a differential response against compounds that trigger apoptosis through their specific interaction with said receptor.

Also encompassed by the present invention is the characterization of biochemical and/or molecular markers that specifically identify an insect cell line undergoing apoptosis. For example, it is possible to isolate specific cDNAs induced during an apoptotic process in specific insect cell lines. Although the death core pathway seems to be phylogenetically conserved (Nagata, S. *Cell* 88:355-365 (1997)), the signal transduction pathway from the receptor to the death core pathway is subject to variation across organisms. Messenger RNAs differentially expressed in insect cells undergoing apoptosis are identified by a number of techniques readily available such as differential display (Bauer, D. *et al.* *Nucleic Acid Res.* 21:4272-4280 (1993)) or subtractive libraries (Sommer, H. *et al.* *EMBO J.* 9:605-613 (1990)). The differentially expressed cDNA-encoded proteins are used as markers for apoptosis in specific insect cell lines.

***Transgenic Plants Comprising a DNA Sequence Encoding a Protein of the VIP3 Class***

A host plant expressing at least one of the sequences of the invention has enhanced resistance to attack by plant pests and is thus better equipped to withstand crop losses associated with such attack. By plant is meant any plant species which can be genetically transformed by methods known in the art. Methods known in the art for plant transformation are discussed below. Host plants include, but are not limited to, those species previously listed as target crops.

**PLANT EXPRESSION CASSETTES**

Methodologies for the construction of plant expression cassettes as well as the introduction of foreign DNA into plants are described in the art. Such expression cassettes may include promoters, terminators, enhancers, leader sequences, introns and other regulatory sequences operably linked to the pesticidal protein coding sequence. It is further recognized that promoters or terminators of the VIP3 genes can be used in expression cassettes.

Toxin genes derived from microorganisms may also differ from plant genes. Plant genes differ from genes found in microorganisms in that their transcribed RNA does not possess defined ribosome binding site sequence adjacent to the initiating methionine. Consequently, microbial genes can be enhanced by the inclusion of a eukaryotic consensus translation initiator at the ATG (Kozak, Cell 44:283-292 (1986)). Clontech (1993/1994 catalog, page 210) has suggested the sequence GTCGACCATGGTC (SEQ ID NO:---) as a consensus translation initiator for the expression of the *E. coli uidA* gene in plants. Further, Joshi (Nucleic Acids Res. 15: 6643-6653 (1987)) has compared many plant sequences adjacent to the ATG and suggests the consensus TAAACAATGGCT (SEQ ID NO:---). In situations where difficulties are encountered in the expression of microbial ORFs in plants, inclusion of one of these sequences at the initiating ATG may improve translation. In such cases the last three nucleotides of the consensus may not be appropriate for inclusion in the modified sequence due to their modification of the second AA residue. Preferred sequences adjacent to the initiating methionine may differ between different plant species. By surveying the sequence of maize genes present in the GenBank/EMBL database it can be discerned which nucleotides adjacent to the ATG should be modified to enhance translation of the toxin gene introduced into maize.

In addition, it has been shown that removal of illegitimate splice sites can enhance expression and stability of introduced genes. Genes cloned from non-plant sources and not optimized for expression in plants may contain motifs which can be recognized in plants as 5' or 3' splice sites. Consequently, the transcription process can be prematurely terminated, generating truncated or deleted mRNA. The toxin genes can be engineered to remove these illegitimate splice sites using multiple techniques. For example, several available methods can be utilized to identify potential splice sites in a DNA sequence. First, potential splice sites may be identified by computer analysis of the DNA sequence. Consensus sequences which identify splice sites are known in the art. See, for example, Goodall, G.J. and Filipowicz, W., EMBO J. 10, 2635-2644 (1991) and Brown, J.W.S., Nucleic Acids Research 14, 9549-9559 (1986). Alternately, one can identify splice sites actually processed by a plant by comparing PCR analysis of cDNA derived from the gene with actual gene products. Shorter than expected products are indicative of splicing. Such smaller products are then cloned and sequenced and the exact location of the splice determined. It is also recognized that a combination of computer searching and PCR analysis can be utilized.

The novel toxin genes of the present invention, either as their native sequence or as optimized synthetic sequences as described above, can be operably fused to a variety of promoters for expression in plants including constitutive, inducible, temporally regulated, developmentally regulated, chemically regulated, tissue-preferred and tissue-specific promoters to prepare recombinant DNA molecules, i.e., chimeric genes. Preferred constitutive promoters include the CaMV 35S and 19S promoters (Fraley *et al.*, U.S. Patent No. 5,352,605, issued October 4, 1994). An additionally preferred promoter is derived from any one of several of the actin genes, which are known to be expressed in most cell types. The promoter expression cassettes described by McElroy *et al.* (Mol. Gen. Genet. 231: 150-160 (1991)) can be easily modified for the expression of the novel toxin gene and are particularly suitable for use in monocotyledonous hosts.

Yet another preferred constitutive promoter is derived from ubiquitin, which is another gene product known to accumulate in many cell types. The ubiquitin promoter has been cloned from several species for use in transgenic plants (e.g. sunflower - Binet *et al.* Plant Science 79: 87-94 (1991), maize - Christensen *et al.* Plant Molec. Biol. 12: 619-632 (1989)). The maize ubiquitin promoter has been developed in transgenic monocot systems and its sequence and vectors constructed for monocot transformation are disclosed in the

patent publication EP 0 342 926. The ubiquitin promoter is suitable for the expression of the novel toxin gene in transgenic plants, especially monocotyledons.

Other promoters useful for the expression of the novel toxin gene in plants, particularly maize, are, for example, tissue-specific or tissue-preferential promoters such as those disclosed in WO 93/07278; chemically inducible promoters disclosed in EP-A 0 332 104, herein incorporated by reference in its entirety.

In addition to promoters, a variety of transcriptional terminators are also available for use in chimeric gene construction using the novel toxin gene of the present invention. Transcriptional terminators are responsible for the termination of transcription beyond the transgene and its correct polyadenylation. Appropriate transcriptional terminators and those which are known to function in plants include the CaMV 35S terminator, the *tm1* terminator, the nopaline synthase terminator, the pea *rbcS* E9 terminator and others known in the art. These can be used in both monocotyledons and dicotyledons.

A number of non-translated leader sequences derived from viruses such as those reported in, for example, WO 96/10083 are also known to enhance expression, and these are particularly effective in dicotyledonous cells. Specifically, leader sequences from Tobacco Mosaic Virus (TMV, the " $\Omega$ -sequence"), Maize Chlorotic Mottle Virus (MCMV), and Alfalfa Mosaic Virus (AMV) have been shown to be effective in enhancing expression (e.g. Gallie *et al.* Nucl. Acids Res. 15: 8693-8711 (1987); Skuzeski *et al.* Plant Molec. Biol. 15: 65-79 (1990)).

Various intron sequences have been shown to enhance expression when added to the 5' regulatory region, particularly in monocotyledonous cells. For example, the introns of the maize *Adh1* gene have been found to significantly enhance the expression of the wild-type gene under its cognate promoter when introduced into maize cells (Callis *et al.*, *Genes Develop.* 1:1183-1200 (1987)).

#### OPTIMIZING *vip3* GENES FOR PLANT EXPRESSION

The pesticidal genes of the invention can be optimized for enhanced expression in plants. See, for example, EPA 0359472; EPA 0385962; WO 91/16432; and, Perlak *et al.*, Proc. Natl. Acad. Sci. 88:3324-3328 (1991). In this manner, the coding sequences can be synthesized which are optimized for plant expression.

In one embodiment of the invention the *vip3A(a)* is made according to the procedure disclosed in U.S. Ser. No. 07/951,715, herein incorporated by reference. In this procedure,



maize preferred codons, i.e., the single codon which most frequently encodes that amino acid in maize, are used. The maize preferred codon for a particular amino acid may be derived, for example, from known gene sequences from maize. Maize codon usage for 28 genes from maize plants is found in Murray *et al.*, Nucleic Acids Research 17: 477-498 (1989), the disclosure of which is incorporated herein by reference. Examples of synthetic sequences made with maize optimized codons are set forth in SEQ ID NO:7 (VIP3A(a)), in SEQ ID NO:19 (VIP3A(b)), and in SEQ ID NO:20 (VIP3A(c)).

In this manner, the nucleotide sequences can be optimized for expression in any plant. It is recognized that all or any part of the gene sequence may be optimized or synthetic. That is, synthetic or partially optimized sequences may also be used.

#### PLANT TRANSFORMATION

The recombinant DNA molecules can be introduced into the plant cell in a number of art-recognized ways. Those skilled in the art will appreciate that the choice of method might depend on the type of plant, i.e. monocot or dicot, targeted for transformation. Suitable methods of transforming plant cells are described, for example, in WO 96/10083 and WO 97/46105, respectively, including microinjection, electroporation, *Agrobacterium*-mediated transformation, direct gene transfer, and ballistic particle acceleration using devices available from Agracetus, Inc., Madison, Wisconsin and Dupont, Inc., Wilmington, Delaware.

An preferred embodiment is the protoplast transformation method for maize as disclosed in European Patent Application EP 0 292 435, as well as in U. S. Patent Number 5,350,689, hereby incorporated by reference in its entirety. One particularly preferred set of embodiments for the introduction of the expression cassettes of the present invention into wheat by microprojectile bombardment can be found in U. S. Patent No. 5,610,042 herein incorporated by reference in its entirety.

Transformation of plants can be undertaken with a single DNA molecule or multiple DNA molecules (*i.e.* co-transformation), and both these techniques are suitable for use with the expression cassettes of the present invention. Numerous transformation vectors are available for plant transformation, and the expression cassettes of this invention can be used in conjunction with any such vectors. The selection of vector will depend upon the preferred transformation technique and the target species for transformation.

Many vectors are available for transformation using *Agrobacterium tumefaciens*. These typically carry at least one T-DNA border sequence and include vectors such as

pBIN19 (Bevan, Nucl. Acids Res. (1984)). In one preferred embodiment, the novel toxin gene of the present invention may be inserted into either of the binary vectors pCIB200 and pCIB2001 for use with *Agrobacterium*, the construction of which is disclosed, for example, in WO 95/33818 (example 35) (see also EP 0 332 104, example 19).

An additional vector useful for *Agrobacterium*-mediated transformation is the binary vector pCIB10 contains a gene encoding kanamycin resistance for selection in plants, T-DNA right and left border sequences and incorporates sequences from the wide host-range plasmid pRK252 allowing it to replicate in both *E. coli* and *Agrobacterium*. Its construction is described by Rothstein *et al.* (Gene 53: 153-161 (1987)). Various derivatives of pCIB10 have been constructed which incorporate the gene for hygromycin B phosphotransferase described by Gritz *et al.* (Gene 25: 179-188 (1983)). These derivatives enable selection of transgenic plant cells on hygromycin only (pCIB743), or hygromycin and kanamycin (pCIB715, pCIB717).

Methods using either a form of direct gene transfer or *Agrobacterium*-mediated transfer usually, but not necessarily, are undertaken with a selectable marker which may provide resistance to an antibiotic (e.g., kanamycin, hygromycin or methotrexate) or a herbicide (e.g., phosphinothricin). The choice of selectable marker for plant transformation is not, however, critical to the invention unless the expression of this resistance and its biochemical activity interferes with the choice of protoxin to toxin conversion chosen for use in creating conditional fertility.

For certain plant species, different antibiotic or herbicide selection markers may be preferred. Selection markers used routinely in transformation include the *nptII* gene which confers resistance to kanamycin and related antibiotics (Messing & Vierra, *Gene* 19: 259-268 (1982); Bevan *et al.*, *Nature* 304:184-187 (1983)), the *bar* gene which confers resistance to the herbicide phosphinothricin (White *et al.*, *Nucl Acids Res* 18:1062 (1990), Spencer *et al.*, *Theor Appl Genet* 79:625-631(1990)), the *hph* gene which confers resistance to the antibiotic hygromycin (Blochinger & Diggelmann, *Mol Cell Biol* 4: 2929-2931), the *dhfr* gene, which confers resistance to methotrexate (Bourouis *et al.*, *EMBO J.* 2: 1099-1104 (1983)), the mannose phosphate isomerase gene, which allows selection on mannose as a carbon source (EP 530 129, WO 94/20627).

One such vector useful for direct gene transfer techniques in combination with selection by the herbicide Basta (or phosphinothricin) is pCIB3064. This vector is based on the plasmid pCIB246, which comprises the CaMV 35S promoter in operational fusion to the

*E. coli* GUS gene and the CaMV 35S transcriptional terminator and is described in the PCT published application WO 93/07278, herein incorporated by reference. Another useful selectable marker is obtained by operably linking a ubiquitin promoter, a synthetic PAT gene and a *nos* terminator. One example of a vector comprising this marker is the plasmid pCIB9804.

An additional transformation vector is pSOG35 which utilizes the *E. coli* gene dihydrofolate reductase (DHFR) as a selectable marker conferring resistance to methotrexate and the construction of which is described, for example, in WO 95/33818 (example 35)

Another transformation vector is the vector pGL2 (Shimamoto *et al.* Nature 338, 274-276 (1989)) which contains the *Streptomyces* hygromycin phosphotransferase gene (hpt) operably linked to the 35S promoter and 35S terminator sequences.

Transgenic plants can also be identified through the use of a scorable marker. Examples of scorable markers useful in the invention are  $\beta$ -glucuronidase, green fluorescent protein, and the *C1* and *B-peru* regulatory genes of the maize anthocyanin pathway. In addition, transgenic plants expressing a VIP3 protein can be identified by screening them for insecticidal activity without the need for either scorable or selectable markers.

Transformation of maize with a DNA sequence encoding a protein of the VIP3 class, but preferably a VIP3A(c) protein according to any of the above methods can be readily achieved by microprojectile bombardment of either immature zygotic embryos or serially-propagatable Type I embryogenic callus.

For transformation using immature zygotic embryos, ears are self-pollinated and immature zygotic embryos are obtained approximately 10 days later. Approximately eight hundred immature zygotic embryos are divided among different target plates containing a medium capable of inducing and supporting the formation of embryogenic callus. The immature zygotic embryos are transferred immediately to the same medium but containing 12% sucrose. After 5 hours, the immature zygotic embryos are bombarded with a plasmid or plasmids using the PDS-1000/He device from BioRad. The plasmid or plasmids comprise a selectable marker, such as a gene conferring resistance to phosphinothricin, or a scorable marker, such as green fluorescent protein, and a gene encoding a protein of the VIP3 class prepared for delivery to and expression in maize according to the above description. The plasmid or plasmids are precipitated onto 1  $\mu$ m gold particles essentially

according to the published procedure from BioRad. The particles are delivered using a burst pressure of 1550 psi of helium. Each target plate is shot twice with the plasmid and gold particle preparation. Since in one embodiment of the invention the plasmid or plasmids comprise a chimeric gene coding for resistance to phosphinothricin this substance could be used to select transformed cells in vitro. If used, the selection agent is applied at 10 mg/L on the day of gene delivery and increased to 40 mg/L after approximately one month. The embryogenic callus so obtained may be regenerated in the presence of the selection agent phosphinothricin if the selectable marker is used. Plants are obtained from the selected embryogenic callus lines. The regenerated plants are assayed for resistance to a susceptible insect. All the plants that are resistant to the insect also express the introduced chimeric gene encoding a protein or proteins of the VIP3 class as evidenced by the detection of VIP3 protein in the plant using an ELISA assay. Plants resistant to the insect and expressing the VIP3 protein are transformed.

For transformation of maize using Type I embryogenic callus, the callus is obtained from immature zygotic embryos using standard culture techniques. For gene delivery, approximately 300 mg of the Type I callus is prepared by either chopping with a scalpel blade or by subculturing 3-5 days prior to gene delivery. Prior to gene delivery, the prepared callus is placed onto semi-solid culture medium again containing 12% sucrose. After approximately 4 hours, the tissue is bombarded using the PDS-1000/He Biolistic device from BioRad. The plasmid or plasmids comprise a selectable marker, such as a gene conferring resistance to phosphinothricin, or a scorable marker, such as green fluorescent protein, and a gene encoding a protein of the VIP3 class prepared for delivery to and expression in maize according to the above description. The plasmids are precipitated onto 1  $\mu$ m gold particles using essentially the standard protocol from BioRad. Approximately 16 hours after gene delivery the callus is transferred to standard culture medium containing 2% sucrose and, if the selectable marker is used, to 1 mg/L phosphinothricin. The callus is subcultured on selection for 8 weeks, after which surviving and growing callus is transferred to standard regeneration medium for the production of plants. The regenerated plants are assayed for resistance to a susceptible insect. All the plants that are resistant to the insect also express the introduced chimeric gene encoding a protein of the VIP3 class as evidenced by the detection of VIP3 protein in the plant using an ELISA assay. Plants resistant to the insect and expressing a protein of the VIP3 class are transformed.

## SUPPLEMENTAL INSECT CONTROL PRINCIPLES

The pesticidal proteins of the invention can be used in combination with Bt  $\delta$ -endotoxins or other insecticidal proteins to increase insect target range. Furthermore, the use of the VIPs of the present invention, but preferably of the VIP3A(c) protein, in combination with Bt  $\delta$ -endotoxins or other insecticidal principles of a distinct nature has particular utility for the prevention and/or management of insect resistance.

The various insecticidal crystal proteins from *Bacillus thuringiensis* have been classified based upon their spectrum of activity and sequence similarity. The classification put forth by Höfte and Whiteley, Microbiol. Rev. 53: 242-255 (1989) placed the then known insecticidal crystal proteins into four major classes. Generally, the major classes are defined by the spectrum of activity, with the Cry1 proteins active against Lepidoptera, Cry2 proteins active against both Lepidoptera and Diptera, Cry3 proteins active against Coleoptera, and Cry4 proteins active against Diptera.

Within each major class, the  $\delta$ -endotoxins are grouped according to sequence similarity. The Cry1 proteins are typically produced as 130-140 kDa protoxin proteins which are proteolytically cleaved to produce active toxin proteins about 60-70 kDa. The active portion of the  $\delta$ -endotoxin resides in the NH<sub>2</sub>-terminal portion of the full-length molecule. Höfte and Whiteley, *supra*, classified the then known Cry1 proteins into six groups, 1Aa, 1Ab, 1Ac, 1B, 1C, and 1D. Since then, proteins classified as Cry1Ea, Cry1Fa, Cry9A, Cry9C and Cry9B have also been characterized.

The spectrum of insecticidal activity of an individual  $\delta$ -endotoxin from *Bacillus thuringiensis* tends to be quite narrow, with a given  $\delta$ -endotoxin being active against only a few insects. Specificity is the result of the efficiency of the various steps involved in producing an active toxin protein and its subsequent ability to interact with the epithelial cells in the insect digestive tract. In one preferred embodiment, expression of VIPs in a transgenic plant is accompanied by the expression of one or more Bt  $\delta$ -endotoxins. Particularly preferred Bt  $\delta$ -endotoxins are those disclosed in U.S. application serial no. 07/951,715, herein incorporated by reference.

It is well known that many  $\delta$ -endotoxin proteins from *Bacillus thuringiensis* are actually expressed as protoxins. These protoxins are solubilized in the alkaline environment of the insect gut and are proteolytically converted by proteases into a toxic core fragment (Höfte and Whiteley, Microbiol. Rev. 53: 242-255 (1989)). For  $\delta$ -endotoxin proteins of the

CryI class, the toxic core fragment is localized in the N-terminal half of the protoxin. It is within the scope of the present invention that genes encoding either the full-length protoxin form or the truncated toxic core fragment of the novel toxin proteins can be used in plant transformation vectors to confer insecticidal properties upon the host plant.

Other insecticidal principles include protease inhibitors (both serine and cysteine types), lectins,  $\alpha$ -amylase, peroxidase and cholesterol oxidase. Other VIP genes, such as *vip1A(a)* and *vip2A(a)* as disclosed in U.S. Ser. No. 08/463,483 and herein incorporated by reference, are also useful in the present invention.

This co-expression of more than one insecticidal principle in the same transgenic plant can be achieved by genetically engineering a plant to contain and express all the genes necessary. Alternatively, a plant, Parent 1, can be genetically engineered for the expression of VIPs. A second plant, Parent 2, can be genetically engineered for the expression of a supplemental insect control principle. By crossing Parent 1 with Parent 2, progeny plants are obtained which express all the genes introduced into Parents 1 and 2.

#### ***Recombinant Microorganisms Comprising Genes and Proteins of the VIP3 Class***

It is recognized that the isolated genes of the present invention which encode a protein of the VIP3 class, but preferably a VIP3A(c) protein, can be transferred into any microbial host and confer their insecticidal properties upon that host. Alternate hosts for the novel genes of the present invention can be selected as suitable for cloning purposes, for purposes of characterizing the form and function of the gene or encoded protein, for use as a fermentation host to increase production of the toxin protein, for purposes of delivering at least one of the toxin proteins more effectively to the target insect pest, or introduction of the novel toxin gene into insect pathogens such as baculovirus (a nuclear polyhedrosis virus, e.g. *Autographica californica*) to improve their effectiveness.

It is envisioned that said alternate host would be applied to the environment or plants or animals for insect control. Microorganism hosts may be selected which are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplana) of one or more crops of interest. These microorganisms are selected so as to be capable of successfully competing in the particular environment with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

Such microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, e.g., *Bacillus*, *Caulobacter*, *Agmenellum*, *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylius*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, e.g., *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Bacillus* spp., *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacteria*, *Rhodopseudomonas spheroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes entrophus*, *Clavibacter xyli* and *Azotobacter vinlandii*; and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces rosues*, *S. odoratus*, *Kluyveromyces veronae*, and *Aureobasidium pollulans*. Of particular interest are the pigmented microorganisms.

Suitable host cells, where the pesticide-containing cells will be treated to prolong the activity of the toxin in the cell when the then treated cell is applied to the environment of the target pest(s), may include either prokaryotes or eukaryotes, normally being limited to those cells which do not produce substances toxic to higher organisms, such as mammals. However, organisms which produce substances toxic to higher organisms could be used, where the toxin is unstable or the level of application sufficiently low as to avoid any possibility of toxicity to a mammalian host. As hosts, of particular interest will be the prokaryotes and the lower eukaryotes, such as fungi. Illustrative prokaryotes, both Gram-negative and -positive, include *Enterobacteriaceae*, such as *Escherichia*, *Erwinia*, *Shigella*, *Salmonella*, and *Proteus*; *Bacillaceae*; *Rhizobiceae*, such as *Rhizobium*; *Spirillaceae*, such as *Photobacterium*, *Zymomonas*, *Serratia*, *Aeromonas*, *Vibrio*, *Desulfovibrio*, *Spirillum*; *Lactobacillaceae*; *Pseudomonadaceae*, such as *Pseudomonas* and *Acetobacter*; *Azotobacteraceae* and *Nitrobacteraceae*. Among eukaryotes are fungi, such as *Phycomycetes* and *Ascomycetes*, which includes yeast, such as *Saccharomyces* and *Schizosaccharomyces*; and *Basidiomycetes* yeast, such as *Rhodotorula*, *Aureobasidium*, *Sporobolomyces*, and the like.

Characteristics of particular interest in selecting a host cell for purposes of production include ease of introducing the protein gene into the host, availability of expression systems, efficiency of expression, stability of the protein in the host, and the presence of auxiliary genetic capabilities. Characteristics of interest for use as a pesticide microcapsule

include protective qualities for the pesticide, such as thick cell walls, pigmentation, and intracellular packaging or formation of inclusion bodies; leaf affinity; lack of mammalian toxicity; attractiveness to pests for ingestion; ease of killing and fixing without damage to the toxin; and the like. Other considerations include ease of formulation and handling, economics, storage stability, and the like.

Host organisms of particular interest include yeast, such as *Rhodotorula sp.*, *Aureobasidium sp.*, *Saccharomyces sp.*, and *Sporobolomyces sp.*; phylloplane organisms such as *Pseudomonas sp.*, *Erwinia sp.*, and *Flavobacterium sp.*; or such other organisms as *Escherichia*, *LactoBacillus sp.*, *Bacillus sp.*, and the like. Specific organisms include *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Saccharomyces cerevisiae*, *Bacillus thuringiensis*, *Escherichia coli*, *Bacillus subtilis*, and the like.

A number of ways are available for introducing a gene expressing the pesticidal protein into the microorganism host under conditions which allow for stable maintenance and expression of the gene. For example, expression cassettes can be constructed which include the DNA constructs of interest operably linked with the transcriptional and translational regulatory signals for expression of the DNA constructs, and a DNA sequence homologous with a sequence in the host organism, whereby integration will occur, and/or a replication system which is functional in the host, whereby integration or stable maintenance will occur.

Transcriptional and translational regulatory signals include but are not limited to promoter, transcriptional initiation start site, operators, activators, enhancers, other regulatory elements, ribosomal binding sites, an initiation codon, termination signals, and the like. See, for example, US Patent 5,039,523; US Patent No. 4,853,331; EPO 0480762A2; Sambrook *et al. supra*; Molecular Cloning, a Laboratory Manual, Maniatis *et al.* (eds) Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982); Advanced Bacterial Genetics, Davis *et al.* (eds.) Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1980); and the references cited therein.

The novel genes or recombinant forms thereof can be transformed into such alternate hosts using a variety of art recognized methods. One such preferred method is electroporation of microbial cells, as described, for example, by the method of Dower (U.S. Patent No. 5,186,800). Another preferred method is that of Schurter *et al.* (Mol. Gen. Genet. 218: 177-181 (1989)), which is also disclosed in U.S. Serial No. 07/353,565 which is incorporated herein in its entirety.



Genes encoding the VIP3 class of proteins can be introduced into microorganisms that multiply on plants (epiphytes) or in plants (endophytes) to deliver proteins of the VIP3 class to potential target pests. Many bacterial species are capable of living in the vascular tissues of plants. Most of these endophytes and epiphytes appear to have little physiological impact on plant growth and productivity.

Root colonizing bacteria, for example, can be isolated from the plant of interest by methods known in the art. Specifically, a *Bacillus cereus* strain which colonizes roots could be isolated from roots of a plant (for example see J. Handelsman, S. Raffel, E. Mester, L. Wunderlich and C. Grau, Appl. Environ. Microbiol. 56:713-718, (1990)). *Vip3* genes can also be introduced into a root colonizing *Bacillus cereus* by standard methods known in the art. Specifically, a gene encoding a protein of the VIP3 class derived from strain AB88 can be introduced into a root colonizing *Bacillus cereus* by means of conjugation using standard methods (J. Gonzalez, B. Brown and B. Carlton, Proc. Natl. Acad. Sci. 79:6951-6955, (1982)).

Also, the novel genes of the invention can be introduced into the root colonizing *Bacillus* by means of electro-transformation. For example, *vip3A(a)* can be cloned into a shuttle vector, for example, pHT3101 (D. Lereclus *et al.*, FEMS Microbiol. Letts., 60:211-218 (1989)). The shuttle vector pHT3101 containing the coding sequence can then be transformed into the root colonizing *Bacillus* by means of electroporation (D. Lereclus *et al.* 1989, FEMS Microbiol. Letts. 60:211-218). It is also possible to use the cotton colonizing *Bacillus megaterium*.

Another example is afforded by the endophyte *Clavibacter xyli*, which is from a genus/species known contain phytopathogenic bacteria which cause plant stunting. This bacterium can grow to very high levels in the vascular system of plants. A  $\delta$ -endotoxin was introduced into this endophyte, which when inoculated into a plant, provided good control of corn borer. Other endophytes are also known.

Expression systems can be designed so that VIP3 proteins are secreted outside the cytoplasm of gram negative bacteria, *E. coli*, for example. Advantages of having VIP3 proteins secreted are (1) it can increase the level of VIP3 protein expressed and (2) can aid in efficient purification of VIP3 protein.

VIP3 proteins can be made to be secreted in *E. coli*, for example, by fusing an appropriate *E. coli* signal peptide to the amino-terminal end of the VIP3 signal peptide or replacing the VIP3 signal peptide with the *E. coli* signal peptide. Signal peptides recognized

by *E. coli* can be found in proteins already known to be secreted in *E. coli*, for example the OmpA protein (J. Ghrayeb, H. Kimura, M. Takahara, Y. Masui and M. Inouye, EMBO J., 3:2437-2442 (1984)). OmpA is a major protein of the *E. coli* outer membrane and thus its signal peptide is thought to be efficient in the translocation process. Also, the OmpA signal peptide does not need to be modified before processing as may be the case for other signal peptides, for example lipoprotein signal peptide (G. Duffaud, P. March and M. Inouye, Methods in Enzymology, 153:492 (1987)).

Specifically, unique BamHI restriction sites can be introduced at the amino-terminal and carboxy-terminal ends of the VIP coding sequences using standard methods known in the art. These BamHI fragments can be cloned, in frame, into the vector pIN-III-ompA1, A2 or A3 (J. Ghrayeb, H. Kimura, M. Takahara, H. Hsiung, Y. Masui and M. Inouye, EMBO J., 3:2437-2442 (1984)) thereby creating ompA:VIP fusion gene which is secreted into the periplasmic space. The other restriction sites in the polylinker of pIN-III-ompA can be eliminated by standard methods known in the art so that the VIP3 amino-terminal amino acid coding sequence is directly after the ompA signal peptide cleavage site. Thus, the secreted VIP3 sequence in *E. coli* would then be identical to the native VIP3 sequence.

When the VIP3 native signal peptide is not needed for proper folding of the mature protein, such signal sequences can be removed and replaced with the ompA signal sequence. Unique BamHI restriction sites can be introduced at the amino-termini of the proprotein coding sequences directly after the signal peptide coding sequences of VIP3 and at the carboxy-termini of VIP3 coding sequence. These BamHI fragments can then be cloned into the pIN-III-ompA vectors as described above.

General methods for employing the strains of the invention in pesticide control or in engineering other organisms as pesticidal agents are known in the art. See, for example US Patent No. 5,039,523 and EP 0480762A2.

VIP3 can be fermented in a bacterial host and the resulting bacteria processed and used as a microbial spray in the same manner that *Bacillus thuringiensis* strains have been used as insecticidal sprays. In the case of a VIP3 which is secreted from *Bacillus*, the secretion signal is removed or mutated using procedures known in the art. Such mutations and/or deletions prevent secretion of the VIP3 protein(s) into the growth medium during the fermentation process. The VIP3 proteins are retained within the cell and the cells are then processed to yield the encapsulated VIP3 protein. Any suitable microorganism can be used for this purpose. *Pseudomonas* has been used to express *Bacillus thuringiensis* endotoxins

as encapsulated proteins and the resulting cells processed and sprayed as an insecticide. (H. Gaertner *et al.* 1993, In Advanced Engineered Pesticides, L. Kim ed.)

Various strains of *Bacillus thuringiensis* are used in this manner. Such *Bt* strains produce endotoxin protein(s) as well as VIP3. Alternatively, such strains can produce only VIP3. A sporulation deficient strain of *Bacillus subtilis* has been shown to produce high levels of the Cry3A endotoxin from *Bacillus thuringiensis* (Agaisse, H. and Lereclus, D., "Expression in *Bacillus subtilis* of the *Bacillus thuringiensis* CryIII<sup>A</sup> toxin gene is not dependent on a sporulation-specific sigma factor and is increased in a *spoOA* mutant", J. Bacteriol., 176:4734-4741 (1994)). A similar *spoOA* mutant can be prepared in *Bacillus thuringiensis* and used to produce encapsulated VIP3 which are not secreted into the medium but are retained within the cell.

Target crops to be protected within the scope of the present invention comprise, e.g., the following species of plants:

cereals (wheat, barley, rye, oats, rice, sorghum and related crops), beet (sugar beet and fodder beet), forage grasses (orchardgrass, fescue, and the like), drupes, pomes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries), leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives, sunflowers, coconuts, castor oil plants, cocoa beans, groundnuts), cucumber plants (cucumber, marrows, melons) fiber plants (cotton, flax, hemp, jute), citrus fruit (oranges, lemons, grapefruit, mandarins), vegetables (spinach, lettuce, asparagus, cabbages and other Brassicaceae, onions, tomatoes, potatoes, paprika), lauraceae (avocados, carrots, cinnamon, camphor), deciduous trees and conifers (e.g. linden-trees, yew-trees, oak-trees, alders, poplars, birch-trees, firs, larches, pines), or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (including composites).

The microorganisms which have been genetically altered to contain the pesticidal gene and protein may be used for protecting agricultural crops and products from pests. In one aspect of the invention, whole, i.e., unlysed, cells of a toxin (pesticide)-producing organism are treated with reagents that prolong the activity of the toxin produced in the cell when the cell is applied to the environment of target pest(s).

Alternatively, the pesticides are produced by introducing a heterologous gene into a cellular host. Expression of the heterologous gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide. These cells are then treated under conditions that prolong the activity of the toxin produced in the cell when the cell is applied to the environment of target pest(s). The resulting product retains the toxicity of the toxin. These naturally encapsulated pesticides may then be formulated in accordance with conventional techniques for application to the environment hosting a target pest, e.g., soil, water, and foliage of plants. See, for example EPA 0192319, and the references cited therein.

The active ingredients of the present invention are normally applied in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession, with other compounds. These compounds can be both fertilizers or micronutrient donors or other preparations that influence plant growth. They can also be selective herbicides, insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired, together with further agriculturally acceptable carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, binders or fertilizers.

Preferred methods of applying an active ingredient of the present invention or an agrochemical composition of the present invention which contains at least one of the pesticidal proteins produced by the bacterial strains of the present invention are leaf application, seed coating and soil application. The number of applications and the rate of application depend on the intensity of infestation by the corresponding pest.

#### ***Entomocidal Compositions Comprising a Recombinant *Bacillus thuringiensis* Strain***

The present invention further provides an entomocidal composition comprising a recombinant *Bacillus thuringiensis* strain containing at least one of the novel toxin genes in recombinant form, or derivatives or mutants thereof, together with an agricultural adjuvant such as a carrier, diluent, surfactant or application-promoting adjuvant. The composition may also contain a further biologically active compound selected from fertilizers, micronutrient donors, plant growth preparations, herbicides, insecticides, fungicides, bactericides, nematocides and molluscicides and mixtures thereof. The composition may comprise from 0.1 to 99% by weight of a recombinant *Bacillus thuringiensis* strain

containing at least one of the novel genes in recombinant form, or the derivatives or mutants thereof, from 1 to 99.9% by weight of a solid or liquid adjuvant, and from 0 to 25% by weight of a surfactant. The recombinant *Bacillus thuringiensis* strain containing at least one of the novel genes in recombinant form, or the composition containing it, may be administered to the plants or crops to be protected together with certain other insecticides or chemicals (1993 Crop Protection Chemicals Reference, Chemical and Pharmaceutical Press, Canada) without loss of potency. It is compatible with most other commonly used agricultural spray materials but should not be used in extremely alkaline spray solutions. It may be administered as a dust, a suspension, a wettable powder or in any other material form suitable for agricultural application.

A recombinant *Bacillus thuringiensis* strain containing at least one of the novel genes in recombinant form is normally applied in the form of entomocidal compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession, with further biologically active compounds. These compounds may be both fertilizers or micronutrient donors or other preparations that influence plant growth. They may also be selective herbicides, insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired together with further carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, binders or fertilizers. The formulations, i.e. the entomocidal compositions, preparations or mixtures containing the recombinant *Bacillus thuringiensis* strain containing the novel gene in recombinant form as an active ingredient or combinations thereof with other active ingredients, and, where appropriate, a solid or liquid adjuvant, are prepared in known manner, e.g., by homogeneously mixing and/or grinding the active ingredients with extenders, e.g., solvents, solid carriers, and in some cases surface-active compounds (surfactants).

Solvents, carriers, surfactants, surface active compounds, etc that are customarily employed in the art of formulation and can be suitably used within the present invention are disclosed, for example, in WO 96/10083.

Another particularly preferred characteristic of an entomocidal composition of the present invention is the persistence of the active ingredient when applied to plants and soil. Possible causes for loss of activity include inactivation by ultra-violet light, heat, leaf

exudates and pH. For example, at high pH, particularly in the presence of reductant,  $\delta$ -endotoxin crystals are solubilized and thus become more accessible to proteolytic inactivation. High leaf pH might also be important, particularly where the leaf surface can be in the range of pH 8-10. Formulation of an entomocidal composition of the present invention can address these problems by either including additives to help prevent loss of the active ingredient or encapsulating the material in such a way that the active ingredient is protected from inactivation. Encapsulation can be accomplished chemically (McGuire and Shasha, 1992) or biologically (Barnes and Cummings, 1986). Chemical encapsulation involves a process in which the active ingredient is coated with a polymer while biological encapsulation involves the expression of the  $\delta$ -endotoxin genes in a microbe. For biological encapsulation, the intact microbe containing the  $\delta$ -endotoxin protein is used as the active ingredient in the formulation. The addition of UV protectants might effectively reduce irradiation damage. Inactivation due to heat could also be controlled by including an appropriate additive.

The entomocidal compositions usually contain 0.1 to 99%, preferably 0.1 to 95%, of a recombinant *Bacillus thuringiensis* strain containing at least one of the novel genes in recombinant form, or combination thereof with other active ingredients, 1 to 99.9% of a solid or liquid adjuvant, and 0 to 25%, preferably 0.1 to 20%, of a surfactant. Whereas commercial products are preferably formulated as concentrates, the end user will normally employ dilute formulations of substantially lower concentration. The entomocidal compositions may also contain further ingredients, such as stabilizers, antifoams, viscosity regulators, binders, tackifiers as well as fertilizers or other active ingredients in order to obtain special effects.

### ***Methods of Controlling Insects***

In view of the above description of the invention, it is apparent that there are several methods by which insects may be controlled using proteins of the VIP3 class as an insecticidal principle, either alone or in combination with supplementary insect control principles such as  $\delta$ -endotoxins. Any method of delivering a VIP3 protein for ingestion by a susceptible insect will result in the control of that insect.

In one embodiment of the invention, plants are transformed with a gene encoding a protein of the VIP3 class. Expression of the protein may occur at any time during growth and development of the plant, depending on the nature of the insect to be controlled. For

example, a protein of the VIP3 class can, according to the invention, be expressed in roots, stems, leaves, seeds, pollen, etc. This provides the advantage of expressing the protein only in those cells or tissues upon which the target insect feeds. Feeding the cells or tissues of a plant expressing VIP3 protein to a susceptible insect will result in the control of that insect. In one embodiment of the invention, a VIP3 protein is expressed in the stem or stalk of a plant in order to control black cutworm. The plants may be grown under either field or greenhouse conditions. Seed containing a VIP3 protein can also be protected against insect damage when in storage.

### EXAMPLES

Examples 16 to 18 on pages 73 to 82 of WO 96/10083 describe the isolation and biological characterization of *Bacillus thuringiensis* strains AB88 and AB424, the purification and characterization of a VIP3A(a) protein, the cloning of *vip3A(a)* and *vip3A(b)* genes, and the identification of new *vip* genes by hybridization. Said Examples are incorporated herein in their entirety by reference. The following examples further describe the materials and methods used in carrying out the invention and the subsequent results. They are offered by way of illustration, and their recitation should not be considered as a limitation of the claimed invention.

#### **Example 1: Presence of *vip3*-like genes and VIP3-like proteins in *Bacillus* isolates**

*Bacillus* isolates other than AB88 have demonstrated insecticidal activity against Lepidopteran larvae when spent culture supernatants were tested. Some isolates which were active against black cutworm were analyzed for the presence of *vip3*-like genes and for the production of VIP3-like proteins.

A standard PCR analysis was used to determine whether the black cutworm-active *Bacillus* isolates contained a *vip3*-like gene. Using the PCR primer pair GW110 (5'-CGA TTA ATG TTG GCC TC-3'; SEQ ID NO:17) and GW111 (5'-CAT TAG CAT CTC CGG ACA CAG-3'; SEQ ID NO:18) it was determined that all of the black cutworm active isolates produced a 728 bp *vip3* gene product which was equal to the size produced by the type strain, AB88. One *Bacillus* isolate, AB51, which was not active against black cutworm, produced the same size *vip3* product. None of the other non-black cutworm active *Bacillus* isolates produced a *vip3* PCR product.

Analysis of VIP3 protein production was done using a standard western blot procedure. Antibodies raised against the VIP3A(a) protein described in the above example were used to detect immunoreactive proteins. Aliquots of cell free culture supernatants from sporulated cultures were run on SDS-PAGE gels using standard methods. Standard western blotting procedures were then carried out to determine the presence of VIP3-like proteins. All of the *Bacillus* isolates which had a 728 bp PCR product and were active against black cutworm produced an 80 kDa protein which was immunoreactive to the VIP3A(a) antibody. The AB51 isolate which had the correct size *vip3* PCR product but was not active against black cutworm produced an immunoreactive protein which was truncated suggesting this may be the reason no biological activity against black cutworm was observed.

**Example 2: Characterization of *Bacillus thuringiensis* Strain AB51 Containing a *vip3*-like Gene**

A *B. thuringiensis* strain, designated AB51, was shown to contain proteins of the VIP3 class by western analysis using rabbit polyclonal anti-Vip3A(a) antibodies. The *vip3*-like gene was cloned into pKS which created pCIB7112. This gene was given the designation *vip3A(c)*. The DNA sequence for *vip3A(c)* is disclosed in SEQ ID NO:5 and the encoded protein sequence is disclosed in SEQ ID NO:6. The VIP3A(c) protein is 746 amino acids long, 43 amino acids shorter than its VIP3A(a) and VIP3A(b) homologues.

**Example 3: Development of Antibodies to VIP3A(a) Protein**

Antiserum against purified Vip3A(a) insecticidal protein was produced in rabbits and goats. For rabbits, nitrocellulose-bound protein (50 µg) was dissolved in DMSO, emulsified with Freund's complete adjuvant (Difco) and injected subcutaneously twice a month for three months. For goats, active soluble pure Vip3A protein (300 µg) was injected intramuscularly twice a month for three month. They were bled 10 days after the second and third injection and the serum was recovered from the blood sample (Harlow, E. and Lane, D. *Antibodies: A Manual Laboratory*, Cold Spring Harbor Lab. Press, NY, 1988). The antisera were then fractionated by affinity chromatography utilizing staphylococcal protein A, and the resulting IgG fraction was further purified by filtering through a column containing immobilized-*E. coli* lysate (Yu, C.G. *et al.* Appl. Environ. Microbiol. 63:532-536 (1997)).



The rabbit and goat antisera were characterized analyzing the Vip3A(a) protein by western blot. Proteins were separated by SDS/PAGE and transferred to nitrocellulose. Nitrocellulose blots were blocked in 20 mM Tris-HCl, pH 7.5/0.15 M NaCl/0.02% NaN<sub>3</sub>/5% nonfat dry milk. Blots were developed by using either rabbit raised or goat-raised anti-Vip3A(a) antibodies at a concentration of 200 ng/ml or 100 ng/ml respectively. Alkaline phosphatase-conjugated goat antirabbit IgG or rabbit antigoaat antiserum were used as secondary antibodies at a concentration of 1 µg/ml (Kirkegaard & Perry Laboratories, Inc.). Bromochloroindolyl-phosphate and nitroblue tetrazolium were used as substrate for the alkaline phosphatase reaction. Both anti-Vip3A(a) antibodies, the rabbit and the goat raised, are polyclonal. The anti-Vip3A(a) antibodies obtained from goat have a higher titer than the ones obtained from rabbits. In the experimental approach, anti-Vip3A(a) antibodies from rabbit should be used at a dilution 1/500 from the original serum (200 ng/ml). By comparison, the anti-Vip3A(a) antibodies obtained from goat can be diluted up to 1/2000 (100 ng/ml) from the original serum. While the rabbit raised antibodies only recognize the N-terminal portion of the Vip3A(a) protein, the antibodies obtained from goats react with epitopes present throughout the full length of the Vip3A(a) protein.

**Example 4: Construction of Plant Expression Cassettes**

Plant expression cassettes consist of promoters that can drive the expression of a coding sequence either constitutively or in a tissue-specific manner, the coding sequences to be expressed and the termination sequences which allow the polyadenylation of the mRNA and its proper translation.

The promoters selected in the DNA constructs of the present invention includes constitutive promoters such as the one from the maize ubiquitin gene (Christensen *et al.* Plant Mol. Biol. 12:619-632, 1989) (pCIB8029, Figure 4; pCIB8055, Figure 5; pCIB9806, Figure 6), and tissue specific promoters such as those from the maize methalothionein-like gene (de Framond, A. FEBS 290:103-106, 1991) (pCIB8030, Figure 7; pCIB8056, pCIB9805) which provides a root-preferred expression, from the maize PEPC gene (Hudspeth, R.L. and Grula, J.W. Plant Mol. Biol. 12:579-589, 1989) (pCIB5535, Figure 8; pCIB9807) which provides a green-tissue specific expression, and from the barley non-specific lipid transfer protein LTP4 (pCIB9819, Figure 9)(Molina, A. and Garcia-Olmedo, F. Plant J. 4:983-991, 1993) which provides a stem-preferred expression. All constructs used in the present invention contain the terminator sequence derived from the 35S CaMV and the intron 9 derived from the maize PEPC gene for enhancing gene expression purposes.

The plasmids pCIB8029, pCIB8055, and pCIB9806 contain the intron#1 of the maize ubiquitin gene placed between the maize ubiquitin promoter and the *vip3A(a)* gene. The construct comprising the encoding sequence of the *vip3A(a)* gene, the intron#9 and the 35S terminator sequence was engineered into the recipient plasmid bearing the different promoters as double digests *BamHI-EcoRI*.

The plant expression cassettes were used as such in the plant transformation experiments, or they were linearised by using restriction enzymes that cut in the  $\text{Amp}^R$  gene of the backbone plasmid. In some experiments, fragments comprising the promoter, gene of interest, intron and terminator were isolated from the rest of the plasmid backbone by restriction digestion and fragment purification. In these cases fragment purification proceeded as follows: 500 ug of DNA is digested with the appropriate enzyme and separated on a 0.8% agarose gel. The fragment of interest is identified, cut out from the gel and purified using a Durapore Millipore filter (0.45 micron). The filtrate containing the fragment is precipitated with sodium acetate and ethanol. The fragment is resuspended in TE and used in transformation experiments.

**Example 5: Insecticidal activity of maize plants expressing VIP3A(a)**

Maize plants expressing VIP3A(a) protein were tested for insecticidal effects on the insect species listed in the table below by the following procedure. One to four 4 cm sections were cut from leaves of transgenic and control maize plants. Each leaf piece was placed on a moistened filter disc in a 50 X 9 mm petri dish. Five neonates of the species being tested were placed on each leaf piece giving a total of 5-20 larvae tested for each plant. The Petri dishes were incubated at 30° C in the dark. Mortality was scored after 48-72 hours. Results are shown in Table 16.

**Table 16.**

Insect species tested	Percent mortality	
	VIP3A(a)	Control
<u>Maize Pests</u>		
Black cutworm ( <i>Agrotis ipsilon</i> )	100	0
Fall armyworm ( <i>Spodoptera frugiperda</i> )	100	0
Sugarcane borer ( <i>Diatrea saccharalis</i> )	100	0
Southwestern corn borer ( <i>Diatraea grandiosella</i> )	100	0
Corn earworm ( <i>Helicoverpa zea</i> )	100	10
Mediterranean corn borer ( <i>Sesamia nonagroides</i> )	100	15

Other Lepidopteran Pests

Beet armyworm ( <i>S. exigua</i> )	100	0
Yellow striped armyworm ( <i>S. ornithogalli</i> )	100	0
Cabbage looper ( <i>Trichoplusia ni</i> )	100	20

Example 6: Expression of vip3A(a) in Maize Plants

Transformation of maize elite Ciba inbred lines CG00526 and 2154 with the Vip3 gene was achieved using particle bombardment of Type I callus tissue. For transformation using Type I embryogenic callus, the callus was obtained from zygotic embryos using standard culture techniques and subcultured 1-2 days prior to bombardment. Callus tissue was prepared for bombardment by placing ~20, 3-5mm diameter pieces arranged in a ring shape onto culture medium containing 12% sucrose. Callus tissue was placed onto this media for four hours prior to bombardment. DNA used for transformation of maize callus was either circular plasmid DNA, linear plasmid DNA, or purified DNA fragments containing the Vip3 gene under control of various plant promoters. In experiments where a selectable agent was used, the gene allowed resistance to phosphinothricin or allowed for growth in the presence of mannose. Plasmids or DNA fragments isolated by filtration were precipitated onto 0.3 um gold particles according to published procedures from BioRad Laboratories, Hercules, CA. Gold particles were delivered using a burst pressure of 650 psi of helium. Each target plate was shot twice with the DNA coated particles. Sixteen to twenty hours after bombardment the CG00526 callus was transferred to standard culture maintenance media. Seven days post-bombardment the tissue was transferred to media containing the selection agent, Basta at a concentration of 100 mg/L. Basta is a commercial formulation of glufosinate ammonium produced by Hoechst. Callus of 2154 was kept on 12% sucrose for 1-7 days after bombardment and transferred to standard culture media containing 20-30 mg/L Basta at day 7. The 2154 and CG00526 callus was subcultured in the presence of 30 or 100 mg/L Basta, respectively, for eight weeks. Tissue surviving selection was subcultured onto lower levels of Basta (5-40 mg/L) for a period of approximately five to ten weeks to allow for tissue bulk-up and then transferred to a standard regeneration media with no selection for the production of plants. Commonly, 12% of the callus pieces bombarded produced transformed callus that survived Basta selection. Individual transformed calli would typically be regenerated to produce 20-30 plants.

Events were generated from experiments where no selection was used. In these experiments the callus was grown for a period of 9-10 weeks on maintenance media prior to

transferring to regeneration media. Event 1337 is an example of a transformed VIP3 event derived from a transformation experiment with no selectable or scorable marker by screening plants for insecticidal activity.

Transformed calli were also generated from experiments where mannose selection was used. In these transformations the phosphomannose isomerase gene under control of the maize ubiquitin promoter of pCIB9818 was bombarded with the Vip3 gene. Mannose at 0.5-1.5% was included in the maintenance media for a period of twelve weeks and not included in the regeneration media.

Transgenic plants were evaluated for VIP3A(a) protein expression by insect bioassay and ELISA assay. Leaf pieces were removed from 2-4 leaf stage plants for evaluation using both black cutworm and fall army worm bioassays. Bioassays were done using ten newly hatched larvae placed in dishes with leaf pieces. Percent mortality was calculated at 72 hours. Tissues from transgenic plants were also assayed by ELISA using standard protocols to quantitate Vip3 protein levels in different plant tissues. Plant tissue was extracted and Table 17 provides representative events generated and their corresponding of insect bioassay results.

Transgenic maize plants were transformed with various plasmids containing the Vip3 gene under control of various promoters such as the maize PEP-carboxylase promoter (PEPC), the maize ubiquitin promoter (Ubi), and the maize metallothionein-like promoter (MTL). The selectable marker gene was the PAT gene under control of the maize ubiquitin promoter in pUBIAC. Representative events listed in Table 17 show the events produced with different plasmids or DNA fragments derived from plasmids. DNA fragments were generated using restriction enzyme digestions and size fractionated using electrophoresis in 0.8% agarose gels. The DNA fragments were excised from the gels, frozen, crushed and purified by filtration through 0.45 micron DuraPore Millipore filters followed by ethanol precipitation. Transformed maize events were generated with circular plasmid DNA of pCIB5535 containing the Vip3 gene under control of the maize PEPC promoter. Events were also transformed with linear plasmid DNA of pCIB5535 and pCIB8029 containing the Vip3 gene under control of the maize ubiquitin promoter. Additional events were produced by bombarding purified DNA restriction enzyme fragments containing just the Vip3 gene with promoter. Fragments corresponding to the Vip3 gene include: a 4906bp EcoRI/HindIII fragment from pCIB5535 with the maize PEPC promoter; a 5142bp KpnI/HindIII fragment from pCIB8030 with the MTL promoter; a 4597bp KpnI/HindIII fragment of pCIB8029 with the maize ubiquitin promoter; a 4818bp HindIII fragment of pCIB8055 with the maize

ubiquitin promoter; a 5364 HindIII fragment of pCIB8056 with the MTL promoter; a 5964 Ascl fragment of pCIB9805 with the MTL promoter; a 5418bp Ascl fragment of pCIB9806 with the maize ubiquitin promoter; and a 5727bp Ascl fragment of pCIB9807 with the maize PEPC promoter.

Table 17

Event No.	Plasmid Used	Chimeric Gene	Mortality (%)	
			Fall Armyworm	Black Cutworm
891	pCIB5535	PEPC: <i>vip3A(a)</i>	100	100
906	pCIB5535 and pCIB8029	PEPC: <i>vip3A(a)</i> and Ubi: <i>vip3A(a)</i>	100	100
946	pCIB5535 and pCIB8030	PEPC: <i>vip3A(a)</i> and MTL: <i>vip3A(a)</i>	100	100

**Example 7: Insecticidal activity of maize plants containing Vip3 and Bt  $\delta$ -endotoxins**

VIP3A(a) has little activity against European corn borer (ECB). To make plants with broad spectrum lepidopteran control, maize plants containing a *vip3A(a)* gene were crossed with maize plants containing a *cry1B*, which is active against ECB. Progeny from the crosses were bioassayed against ECB and fall armyworm (FAW) as described in Example 1. Results are shown in Table 18. Approximately 34 % of the progeny were not active against either species, 15.4 % were active only on ECB, 23.1 % were active only on FAW and 27.9 % were active against both species. Plants active against both species contained both VIP3A(a) and Cry1B protein. Similar results are obtained using other Bt  $\delta$ -endotoxins, particularly Cry1Ab or Cry9C.

Table 18

Cross	% ECB active	% FAW active	% ECB & FAW active	% not active
VIP3A(a) X Cry1B	15.4	23.1	27.9	34.6

**Example 8: VIP3A(a) Lyses the Midgut Epithelial Cells of Susceptible Insects**

**Feeding and gut clearance studies.** The temporal sequence of symptoms following the ingestion of VIP3A(a)-containing diet by second-instar black cutworm (BCW) larvae, a susceptible insect, were recorded from the time of initial administration until larval death. Larvae exposed to control diet showed active feeding followed by uninterrupted gut parastalsis. In contrast, the addition of VIP3A(a) protein in the diet had a significant effect on feeding behavior. When added at concentrations as low as 4 ng per cm<sup>2</sup>, the larvae fed on and off during periods of 10-20 min. The presence of blue color in their guts indicated feeding but the clearance of the gut content was dramatically affected as judged by the decreased number of frass. With 4 ng of VIP3A(a) per cm<sup>2</sup> added to the diet, larval development was significantly impaired after a 48 h incubation period but no mortality was observed. At concentrations of 40 ng of Vip3A(a) per cm<sup>2</sup>, the larvae suffered gut paralysis upon ingestion of minute amounts of diet and no frass could be seen indicating an almost complete lack of gut clearance. Under this condition, ca. 50% mortality was recorded after 48 hr. When concentrations higher than 40 ng of VIP3A(a) per cm<sup>2</sup> were used, the larvae were moribund after only a few bites, with no frass and mortality rates approaching 100%. When similar experiments were conducted with fall armyworm, also a susceptible insect, similar behavioral patterns were observed. In contrast, European corn borer did not alter its feeding behavior when VIP3A(a) protein was added to the diet even at concentrations as high as 400 ng of VIP3A(a) per cm<sup>2</sup>.

**Histological observations of the effects of the Vip3A(a) protein.** Histopathological observations on the effects of the VIP3A(a) protein on BCW were conducted on second and third instar larvae which had been fed a diet containing VIP3A(a). Analysis of BCW gut cross-sections showed extensive damage to the midgut epithelium indicating that the midgut tissue is a primary site of action of the Vip3A(a) protein. No damage was discernible in the foregut and hindgut. Midgut epithelial cells from untreated larvae were closely associated with one another showing no evidence of damage. Sections from larvae that had been fed for 24 h with diet containing Vip3A(a) showed that distal ends of the epithelium columnar cells had become distended and bulbous. Although the goblet cells exhibited some morphological alterations, they did not show signs of damage at this stage. Degeneration of the epithelium columnar cells continued such that, after 48 h of ingesting Vip3A(a)-containing diet, the lumen was filled with debris of disrupted cells. The goblet cells also exhibited signs of damage after 48 h, but both types of cells were still attached to the basement membrane. Black cutworm larvae were dead at 72h and desquamation of the

epithelial layer was complete. While a similar histopathology was observed for fall armyworm, European corn borer did not exhibit any tissue damage under similar experimental conditions.

***In vivo* immunolocalization of the Vip3A(a) protein.** Third instar larvae of black cutworm and European corn borer fed on artificial diet supplemented with 100-200 ng of VIP3A(a) per cm<sup>2</sup> were used for immunocytochemical characterization of the VIP3A(a) binding to midgut sections. The bound VIP3A(a) was visualized using rabbit anti-VIP3A(a) antibodies previously purified through protein A sepharose and *E. coli* immobilized columns (Yu, C.G. *et al.* Appl. Environ. Microbiol. 63:532-536, 1997). VIP3A(a) binding was detected in midgut epithelium of black cutworm, while showing no binding to European corn borer midguts. Midgut sections from black cutworm larvae fed with control diet showed no VIP3A(a) binding. The VIP3A(a) binding seems to be specifically associated to the apical microvilli and it is mostly associated to the columnar cells, with no detectable signal in the goblet cells.

**Example 9: VIP3A(a) and VIP3A(b) Induce Apoptosis in Insect Cells**

VIP3A(a) and VIP3A(b) were shown to be a apoptosis inducing protein arose by the characterization of its insecticidal effects towards an insect cell line (Sf-9) derived from *Spodoptera frugiperda*, an insect susceptible to VIP3A(a). VIP3A(a) showed insecticidal activity towards the insect cell line when kept present throughout the experiment. When SF-9 insect cells are transiently exposed to VIP3A(a) and VIP3A(b), their cell viability was significantly reduced even with exposure times as short as 5 min. Once the incubation time exceeded 10 min, the effects of the VIP3A(a) and VIP3A(b) on insect cell viability over a period of 6 hours were maximal showing a reduction of 90% in cell viability. The cytological changes occurring in SF-9 cells transiently exposed to VIP3A(a) were monitored by microscopy. Small protrusions appeared on the surface of the treated cells some time between 10 and 15 min after their exposure to the VIP3A(a) protein. At this stage, the mitochondria of the cells remained functionally intact as revealed by staining with rhodamine 123, a dye that accumulates in mitochondria with active membrane potential (Johnson, L.V. *et al.* Proc. Natl. Acad. Sci. USA 77:990-994, 1980). These protrusions eventually disappeared and the cells entered a phase of profuse vacuolization lasting an additional 30 to 60 min. During the final stages, the insect cells are seen to swell before disintegration. For an individual cell, the entire process required 1 to 2 hours. All these cellular events are consistent with previous studies on cells undergoing apoptosis

particularly considering that programmed cell death during metamorphosis of certain insects is accompanied by cellular vacuolization and swelling (Schwartz, L.M. *et al.* Proc. Natl. Acad. Sci. USA 90:980-984 (1993)).

Recent studies have shown that the distribution of phospholipids in the plasma membrane is affected in very early stages of animal cells undergoing apoptosis (Martin, S. J., *et al.* J. Exp. Med.:182, 1545-1556, 1995) particularly the externalization of the phosphatidylserine (PS). This process can be visualized by using Annexin V, an anticoagulant protein with high affinity for phosphatidylserine (PS). When VIP3A(a)-treated SF-9 cells were incubated with Annexin V, an externalization of PS was revealed in insect cell membranes as early as 5-10 min after the exposure to VIP3A(a) probably marking the onset of apoptosis.

One of the key molecular events that is the hallmark of apoptosis is endonucleolysis resulting in a double strand DNA break freeing oligonucleosome-sized fragments of 200 base pair and multiples. We examined the occurrence of endonucleolysis in SF-9 cells treated with VIP3A(a) using an *in situ* detection method and analysing the DNA by agarose gel electrophoresis. Based on the ability of the Klenow enzyme to incorporate modified nucleotides using the DNA ends generated by DNA fragmentation, SF-9 insect cells showed signs of endonucleolysis as early as 30 min upon their exposure to the VIP3A(a) protein. This stage will coincide with the appearance of membrane-bound subcellular apoptotic bodies visualized in the microscopical observations. These early indications of endonucleolytic activity were confirmed by the detection of DNA fragments in agarose gels characteristic of a chromatin ladder slightly latter in the process. These results corroborated the indications obtained from cytological observations, that the SF-9 cells initiate an apoptotic-type of programmed cell death upon their exposure to the VIP3A(a) protein.

The VIP3A(a) and VIP3A(b) proteins were discovered on the basis of their insecticidal properties against some lepidopteran insects. Therefore, we were interested in knowing whether the VIP3A(a) protein would induce an apoptotic pathway in gut cells of susceptible insects upon its ingestion and thus, it could exert its insecticidal properties by triggering an active process of cell death. Histological and histochemical studies have shown that the VIP3A(a) protein specifically targets the columnar cells of the midgut epithelium of susceptible insects provoking cell changes characterized by membrane protrusions and extensive vacuolization leading to cell death. These cytological changes induced by VIP3A(a) in insect gut cells resemble those described above for SF-9 cells. We then



examined whether midgut epithelium cells of susceptible insects undergo endonucleolysis upon ingesting diet containing VIP3A(a) by *in situ* detection (Cuvillier, O., *et al.* Nature 381:800-803 (1996)) of DNA fragmentation. When sections of midgut tissue from black cutworm larvae fed with diet either containing VIP3A(a) or control diet, nuclei staining indicative of DNA fragmentation was only detectable in the columnar cells of the midgut epithelium exposed to the VIP3A(a) protein. This result indicates that the VIP3A(a) protein induces an endonucleolysis process in the midgut epithelium cells concurrently with the cytological changes reported previously. It is our conclusion that the VIP3A(a) protein likely exerts its insecticidal properties by activating an apoptosis-type of programmed cell death of the midgut epithelium cells of susceptible insects.

**Example 10: Isolation of the Receptor for VIP3A(a) from black cutworm**

Black cutworm is sensitive to VIP3A(a) and therefore this insect was used for the isolation of the VIP3A(a) receptor. Midgut of third instar black cutworm larvae were collected by dissection and immediately frozen in liquid nitrogen. One gram of midgut tissue was used to isolate mRNA by following the protocol described in the two-hybrid cDNA library construction kit provided by Clontech (1997). Ten micrograms of poly A<sup>+</sup> RNA were used as starting material. In first strand synthesis, both random and lock-docking oligo(dT)<sub>25</sub>d(A/C/G) primers are used in separate synthesis with MML reverse transcriptase. The second strand cDNA was achieved by an optimal ratio of DNA polymerase to Rnase H activity in the second-strand enzyme cocktail. The newly synthesized double stranded cDNA is then ligated to *EcoRI*-*NotI*-*SalI* adaptors. The cDNAs were ligated into pGAD10 (Vijaychander, S. *et al.* CLONTECHniques IX-3:8-10, 1994) which provides the activation domain. The *vip3A(a)* gene was engineered into the polylinker site of the plasmid pGBT9 in frame with the GAL4-DNA binding domain (Bartel, P.L. *et al.* Cellular Interactions in Development: A Practical Approach, pp. 153-179, Oxford University Press, 1993). The recombinant pGBT9-*vip3A(a)* was transformed into the yeast strain GGY1::171 (Gill, G. and Ptashne, M. Cell51:121-126, 1987) by electroporation (Estruch, J.J. *et al.* BioTechniques 16:610-612, 1994). The transformed yeast was selected in minimal media without tryptophan (Bartel, P.L. *et al.* Cellular Interactions in Development: A Practical Approach, pp. 153-179, Oxford University Press, 1993). The expression of the VIP3A(a) protein in the recombinant yeast was confirmed by western analysis. The yeast strain GGY1::171- VIP3A(a) was transformed with the black cutworm cDNA library represented in pGAD10. GGY1::171 possess the HIS3

marker under the control of GAL4 recognition sites. The HIS3 gene allows a positive growth selection for clones that are transformed by two interacting hybrid constructs. After plating more than 200,000 recombinant clones, only one was able to grow in minimal media without histidine. The plasmid DNA of the positive yeast colony was isolated by the yeast lysis buffer method (Kaiser, P. and Auer, B. *BioTechniques* 14:552 (1993)) and electroporated into *E. coli*. The insert containing the cDNA was subcloned into the *EcoRI* site of the pBluescript (Stratagene) and sequenced by the dideoxy termination method of Sanger *et al.*, *Proc. Natl. Acad. Sci. USA*, 74: 5463-5467 (1977), using PRISM Ready Reaction Dye Deoxy Terminator Cycle Sequencing Kits and PRISM Sequenase® Terminator Double-Stranded DNA Sequencing Kit and analysed on an ABI 373 automatic sequencer.

An alternative approach to identify clones encoding for protein(s) that interact with Vip3A consisted of plating yeast transformed with the black cutworm cDNA library represented in pGAD10. After transferring said transformed yeast population to nitrocellulose filters, they were screened with a biotin-labelled Vip3A protein followed by an incubation with a solution containing streptavidin coupled to alkaline phosphatase (AP). After extensive washes, the clone or clones expressing a protein with the ability to bind to Vip3A were visualized by using the substrate 5-bromo-4-chloro-3-indolyl phosphate (BCIP) in combination with nitro blue tetrazolium (NBT). AP catalyzes the formation of insoluble precipitates which developed in dark spots on the nitrocellulose filters. The experimental procedures are described in detail by Sambrook, J. *et al.* in *Molecular Cloning: A Laboratory Manual*, pp. 12-21-12.24, Cold Spring Harbor Laboratory, 1989. After re-growing the yeast plates overnight, the developed spots in the filters were matched to yeast colonies in the original plates. These colonies were grown, their plasmids isolated, and their inserts characterized as described above.

**Example 11: Insect Cells Transformed with the Gene for the Receptor Exhibit Apoptosis when Exposed to the VIP3A(a) protein**

The receptor in black cutworm midgut cells for the VIP3A(a) protein was cloned into the *XhoI-BamHI* site of the Smart 2 cosmid vector (Speck, M. *et al* *Gene* 64: 173-177 (1988)), and the recombinant construct was used to transform the Schneider 2 (S2) *Drosophila* cell line using the calcium phosphate co-precipitation method (Clem, R.J. and Miller, L. K. *Mol. Cel. Biol.* 14: 5212-5222 (1994)). Smart 2 carries the selectable marker *tet* (tetracycline) for bacterial transformation and the *neo* (neomycin) for *Drosophila* cell transformation. The neo selectable marker is expressed under the control of the *Drosophila*

hsp70 promoter. The transformed S2 cells were selected in S2 *Drosophila* medium supplemented with 10% of Fetal Seroalbumin and with G418 (1mg/ml) at 30°C (see GIBCO catalogue 1997). Several stably transformed S2 cell lines were established after 45 days of selection in the medium described above.

The sensitivity of the S2 transformed cells to the VIP3A(a) was tested by adding VIP3A(a) protein (at a final concentration of 1,7 micrograms per ml) to the media containing the transformed S2 cells that have been previously heat shocked at 42° C 30 min. The induction of apoptosis in transformed S2 cells was confirmed by both microscopical observations and by the TACS Kit, and *in situ* Apoptosis Detection kit (for detailed description, see Trevigen catalogue 1996).

**Example 12: Isolating Homologues to the Receptor from Other Insects**

The cells of the midgut epithelium of black cutworm larvae possess a receptor that is recognized by the VIP3A(a) protein. Receptors from other insects known to be susceptible to VIP3A(a) are isolated by identifying the DNA sequences in Southern analysis. DNA is prepared, enzyme restricted, run in agarose gels and blotted onto nitrocellulose and/or nylon filters. These filters are probed with the cDNA encoding the receptor from black cutworm using low stringency conditions of hybridization and washing. Genes with a similarity to the black cutworm receptor to VIP3A(a) lower than 50% were identified. The Southern analysis can also be probed against partial sequences of the cDNA which encode specific domains such as death domain or EGF-like motifs with the intention of isolating genes that contain similar domains even though they are functionally different to the black cutworm receptor to VIP3A(a).

The isolation of homologues to the black cutworm receptor to VIP3A(a) is be accomplished by the two hybrid system described in Fields, S. and Song, O.-K. Nature 340:245-246 (1989). Isolated mRNA is obtained from an organisms of interest, synthesize cDNAs and clone them into pGAD10 or equivalent plasmids. The cDNA library is co-transformed with the pGB9-bearing the *vip3A(a)* gene (or homologues of this gene) and rescued putative receptors in yeast by means of activating a marker based upon protein-protein interaction between the VIP3A(a) protein (or homologues) and the putative receptor.

Homologues to the black cutworm receptor to VIP3A(a) are isolated by expressing cDNAs libraries isolated from organisms of interests, cloned into appropriate expression vectors and transformed into host cells such as yeast or insect cells which are known not to

have the ability to bind and/or be sensitive to VIP3A(a). The transformed cells are screened based on their gained property of binding VIP3A(a) or undergoing apoptotic responses when incubated with VIP3A(a). In this case, the protein VIP3A(a) is used as probe and its binding will be monitored either by antibodies against VIP3A(a) or by labels such as biotin attached to VIP3A(a).

**Example 13: Screening for Novel Compounds that Induce Apoptosis in Insect Cells**

Model cell lines for different orders of insects (some examples include Sf-9 cells for lepidopteran, Colorado potato beetle for coleopterans, S2 from *Drosophila* for dipterans) is used to screen for novel compounds whose mode of action is induction of apoptosis. The cells are grown in multi-well plates which are used for a high-throughput assay screening for thousands of compounds (both of large and small molecular weight). The compound(s) are added as single component or as mixtures. Compound(s) inducing apoptosis are identified as follows: 1) membrane protrusions are visible in the cell membrane, 2) a reorganization of the phosphatidylserine containing membrane lipids is detectable by using specific proteins with high affinity for phosphatidylserine such as Annexin-V linked to a visual marker, 3) cytoplasmic blebbing is visible in the cell cytoplasm, 4) active mitochondria can be visualized by using vital dyes such as rhodamine 123 that accumulate in mitochondria, 5) DNA fragmentation is detected either by DNA analysis in agarose gels, by ELISA detection of nucleosomal release or by *in vivo* detection of DNA nicking. All these cytological and molecular features are indicative of apoptosis.

The black cutworm receptor to VIP3A(a) is transformed into S2 cell line. Therefore, isogenic S2 lines are available with and without the said receptor. These cell lines are used to screen compounds that provide a differential response due to the presence of the said receptor. Transformed S2 cells undergoing apoptosis upon exposure to certain compounds are identified as indicated above. The differential response of the transformed versus the non-transformed cell is indicative that the action of the compound is mediated by the cloned receptor. Similar approaches are undertaken with insect cells transformed with receptors homologue to the black cutworm receptor to VIP3A(a).

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: NOVARTIS AG
- (B) STREET: Schwarzwaldallee 215
- (C) CITY: Basel
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE (ZIP): 4058
- (G) TELEPHONE: +4161 324 11 11
- (H) TELEFAX: + 4161 322 75 32

.....(ii) TITLE OF INVENTION: A Novel Class of Proteins for the  
Control of Plant Pests

(iii) NUMBER OF SEQUENCES: 20

## .....(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2378 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 9..2375
- (D) OTHER INFORMATION: /note= "Native DNA sequence  
encoding VIP3A(a) protein from AB88 as contained in pCIB7104"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGATGAAC ATG AAC AAG AAT AAT ACT AAA TTA AGC ACA AGA GCC TTA CCA  
Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro

50

1	5	10	
AGT TTT ATT GAT TAT TTT AAT GGC ATT TAT GGA TTT GCC ACT GGT ATC Ser Phe Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile 15 20 25 30			98
AAA GAC ATT ATG AAC ATG ATT TTT AAA ACG GAT ACA GGT GGT GAT CTA Lys Asp Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu 35 40 45			146
ACC CTA GAC GAA ATT TTA AAG AAT CAG CAG TTA CTA AAT GAT ATT TCT Thr Leu Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser 50 55 60			194
GGT AAA TTG GAT GGG GTG AAT GGA AGC TTA AAT GAT CTT ATC GCA CAG Gly Lys Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln 65 70 75			242
GGA AAC TTA AAT ACA GAA TTA TCT AAG GAA ATA TTA AAA ATT GCA AAT Gly Asn Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn 80 85 90			290
GAA CAA AAT CAA GTT TTA AAT GAT GTT AAT AAC AAA CTC GAT GCG ATA Glu Gln Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile 95 100 105 110			338
AAT ACG ATG CTT CGG GTA TAT CTA CCT AAA ATT ACC TCT ATG TTG AGT Asn Thr Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser 115 120 125			386
GAT GTA ATG AAA CAA AAT TAT GCG CTA AGT CTG CAA ATA GAA TAC TTA Asp Val Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu 130 135 140			434
AGT AAA CAA TTG CAA GAG ATT TCT GAT AAG TTG GAT ATT ATT AAT GTA Ser Lys Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val 145 150 155			482
AAT GTA CTT ATT AAC TCT ACA CTT ACT GAA ATT ACA CCT GCG TAT CAA Asn Val Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln 160 165 170			530
AGG ATT AAA TAT GTG AAC GAA AAA TTT GAG GAA TTA ACT TTT GCT ACA Arg Ile Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr 175 180 185 190			578
GAA ACT AGT TCA AAA GTA AAA AAG GAT GGC TCT CCT GCA GAT ATT CTT Glu Thr Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu 195 200 205			626
GAT GAG TTA ACT GAG TTA ACT GAA CTA GCG AAA AGT GTA ACA AAA AAT Asp Glu Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn 210 215 220			674
GAT GTG GAT GGT TTT GAA TTT TAC CTT AAT ACA TTC CAC GAT GTA ATG Asp Val Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met 225 230 235			722
GTA GGA AAT AAT TTA TTC GGG CGT TCA GCT TTA AAA ACT GCA TCG GAA Val Gly Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu 240 245 250			770
TTA ATT ACT AAA GAA AAT GTG AAA ACA AGT GGC AGT GAG GTC GGA AAT Leu Ile Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn 255 260 265 270			818

GTT TAT AAC TTC TTA ATT GTA TTA ACA GCT CTG CAA GCC CAA GCT TTT	866
Val Tyr Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Gln Ala Phe	
275 280 285	
CTT ACT TTA ACA ACA TGC CGA AAA TTA TTA GGC TTA GCA GAT ATT GAT	914
Leu Thr Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp	
290 295 300	
TAT ACT TCT ATT ATG AAT GAA CAT TTA AAT AAG GAA AAA GAG GAA TTT	962
Tyr Thr Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe	
305 310 315	
AGA GTA AAC ATC CTC CCT ACA CTT TCT AAT ACT TTT TCT AAT CCT AAT	1010
Arg Val Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn	
320 325 330	
TAT GCA AAA GTT AAA GGA AGT GAT GAA GAT GCA AAG ATG ATT GTG GAA	1058
Tyr Ala Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu	
335 340 345 350	
GCT AAA CCA GGA CAT GCA TTG ATT GGG TTT GAA ATT AGT AAT GAT TCA	1106
Ala Lys Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser	
355 360 365	
ATT ACA GTA TTA AAA GTA TAT GAG GCT AAG CTA AAA CAA AAT TAT CAA	1154
Ile Thr Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln	
370 375 380	
GTC GAT AAG GAT TCC TTA TCG GAA GTT ATT TAT GGT GAT ATG GAT AAA	1202
Val Asp Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys	
385 390 395	
TTA TTG TGC CCA GAT CAA TCT GAA CAA ATC TAT TAT ACA AAT AAC ATA	1250
Leu Leu Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile	
400 405 410	
GTA TTT CCA AAT GAA TAT GTA ATT ACT AAA ATT GAT TTC ACT AAA AAA	1298
Val Phe Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys	
415 420 425 430	
ATG AAA ACT TTA AGA TAT GAG GTA ACA GCG AAT TTT TAT GAT TCT TCT	1346
Met Lys Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser	
435 440 445	
ACA GGA GAA ATT GAC TTA AAT AAG AAA AAA GTA GAA TCA AGT GAA GCG	1394
Thr Gly Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala	
450 455 460	
GAG TAT AGA ACG TTA AGT GCT AAT GAT GAT GGG GTG TAT ATG CCG TTA	1442
Glu Tyr Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu	
465 470 475	
GGT GTC ATC AGT GAA ACA TTT TTG ACT CCG ATT AAT GGG TTT GGC CTC	1490
Gly Val Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu	
480 485 490	
CAA GCT GAT GAA AAT TCA AGA TTA ATT ACT TTA ACA TGT AAA TCA TAT	1538
Gln Ala Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr	
495 500 505 510	
TTA AGA GAA CTA CTG CTA GCA ACA GAC TTA AGC AAT AAA GAA ACT AAA	1586
Leu Arg Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys	
515 520 525	



TTG ATC GTC CCG CCA AGT GGT TTT ATT AGC AAT ATT GTA GAG AAC GGG Leu Ile Val Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly 530 535 540	1634
TCC ATA GAA GAG GAC AAT TTA GAG CCG TGG AAA GCA AAT AAT AAG AAT Ser Ile Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn 545 550 555	1682
GCG TAT GTA GAT CAT ACA GGC GGA GTG AAT GGA ACT AAA GCT TTA TAT Ala Tyr Val Asp His Thr Gly Val Asn Gly Thr Lys Ala Leu Tyr 560 565 570	1730
GTT CAT AAG GAC GGA GGA ATT TCA CAA TTT ATT GGA GAT AAG TTA AAA Val His Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys 575 580 585 590	1778
CCG AAA ACT GAG TAT GTA ATC CAA TAT ACT GTT AAA GGA AAA CCT TCT Pro Lys Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser 595 600 605	1826
ATT CAT TTA AAA GAT GAA AAT ACT GGA TAT ATT CAT TAT GAA GAT ACA Ile His Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr 610 615 620	1874
AAT AAT AAT TTA GAA GAT TAT CAA ACT ATT AAT AAA CGT TTT ACT ACA Asn Asn Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr 625 630 635	1922
GGA ACT GAT TTA AAG GGA GTG TAT TTA ATT TTA AAA AGT CAA AAT GGA Gly Thr Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly 640 645 650	1970
GAT GAA GCT TGG GGA GAT AAC TTT ATT ATT TTG GAA ATT AGT CCT TCT Asp Glu Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser 655 660 665 670	2018
GAA AAG TTA TTA AGT CCA GAA TTA ATT AAT ACA AAT AAT TGG ACG AGT Glu Lys Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser 675 680 685	2066
ACG GGA TCA ACT AAT ATT AGC GGT AAT ACA CTC ACT CTT TAT CAG GGA Thr Gly Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly 690 695 700	2114
GGA CGA GGG ATT CTA AAA CAA AAC CTT CAA TTA GAT AGT TTT TCA ACT Gly Arg Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr 705 710 715	2162
TAT AGA GTG TAT TTT TCT GTG TCC GGA GAT GCT AAT GTA AGG ATT AGA Tyr Arg Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg 720 725 730	2210
AAT TCT AGG GAA GTG TTA TTT GAA AAA AGA TAT ATG AGC GGT GCT AAA Asn Ser Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys 735 740 745 750	2258
GAT GTT TCT GAA ATG TTC ACT ACA AAA TTT GAG AAA GAT AAC TTT TAT Asp Val Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr 755 760 765	2306
ATA GAG CTT TCT CAA GGG AAT AAT TTA TAT GGT GGT CCT ATT GTA CAT Ile Glu Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His 770 775 780	2354
TTT TAC GAT GTC TCT ATT AAG TAA	2378

Phe Tyr Asp Val Ser Ile Lys  
785

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Asn	Lys	Asn	Asn	Thr	Lys	Leu	Ser	Thr	Arg	Ala	Leu	Pro	Ser	Phe	1	5	10	15
Ile	Asp	Tyr	Phe	Asn	Gly	Ile	Tyr	Gly	Phe	Ala	Thr	Gly	Ile	Lys	Asp	20	25	30	
Ile	Met	Asn	Met	Ile	Phe	Lys	Thr	Asp	Thr	Gly	Gly	Asp	Leu	Thr	Leu	35	40	45	
Asp	Glu	Ile	Leu	Lys	Asn	Gln	Gln	Leu	Leu	Asn	Asp	Ile	Ser	Gly	Lys	50	55	60	
Leu	Asp	Gly	Val	Asn	Gly	Ser	Leu	Asn	Asp	Leu	Ile	Ala	Gln	Gly	Asn	65	70	75	80
Leu	Asn	Thr	Glu	Leu	Ser	Lys	Glu	Ile	Leu	Lys	Ile	Ala	Asn	Glu	Gln	85	90	95	
Asn	Gln	Val	Leu	Asn	Asp	Val	Asn	Asn	Lys	Leu	Asp	Ala	Ile	Asn	Thr	100	105	110	
Met	Leu	Arg	Val	Tyr	Leu	Pro	Lys	Ile	Thr	Ser	Met	Leu	Ser	Asp	Val	115	120	125	
Met	Lys	Gln	Asn	Tyr	Ala	Leu	Ser	Leu	Gln	Ile	Glu	Tyr	Leu	Ser	Lys	130	135	140	
Gln	Leu	Gln	Glu	Ile	Ser	Asp	Lys	Leu	Asp	Ile	Ile	Asn	Val	Asn	Val	145	150	155	160
Leu	Ile	Asn	Ser	Thr	Leu	Thr	Glu	Ile	Thr	Pro	Ala	Tyr	Gln	Arg	Ile	165	170	175	
Lys	Tyr	Val	Asn	Glu	Lys	Phe	Glu	Glu	Leu	Thr	Phe	Ala	Thr	Glu	Thr	180	185	190	
Ser	Ser	Lys	Val	Lys	Lys	Asp	Gly	Ser	Pro	Ala	Asp	Ile	Leu	Asp	Glu	195	200	205	
Leu	Thr	Glu	Leu	Thr	Glu	Leu	Ala	Lys	Ser	Val	Thr	Lys	Asn	Asp	Val	210	215	220	
Asp	Gly	Phe	Glu	Phe	Tyr	Leu	Asn	Thr	Phe	His	Asp	Val	Met	Val	Gly	225	230	235	240
Asn	Asn	Leu	Phe	Gly	Arg	Ser	Ala	Leu	Lys	Thr	Ala	Ser	Glu	Leu	Ile	245	250	255	
Thr	Lys	Glu	Asn	Val	Lys	Thr	Ser	Gly	Ser	Glu	Val	Gly	Asn	Val	Tyr	260	265	270	

Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Gln Ala Phe Leu Thr  
 275 280 285  
 Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr  
 290 295 300  
 Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val  
 305 310 315 320  
 Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala  
 325 330 335  
 Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys  
 340 345 350  
 Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr  
 355 360 365  
 Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp  
 370 375 380  
 Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu  
 385 390 395 400  
 Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe  
 405 410 415  
 Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys  
 420 425 430  
 Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly  
 435 440 445  
 Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr  
 450 455 460  
 Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val  
 465 470 475 480  
 Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala  
 485 490 495  
 Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg  
 500 505 510  
 Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile  
 515 520 525  
 Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile  
 530 535 540  
 Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr  
 545 550 555 560  
 Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His  
 565 570 575  
 Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys  
 580 585 590  
 Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His  
 595 600 605  
 Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn

- 62 -

610	615	620
Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr		
625	630	635 640
Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu		
	645	650 655
Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys		
	660	665 670
Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly		
	675	680 685
Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg		
	690	695 700
Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg		
	705	710 715 720
Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser		
	725	730 735
Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val		
	740	745 750
Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr Ile Glu		
	755	760 765
Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His Phe Tyr		
	770	775 780
Asp Val Ser Ile Lys		
785		

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2612 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 118..2484
- (D) OTHER INFORMATION: /note= "Native DNA sequence encoding VIP3A(b) from AB424"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATTGAAATTG ATAAAAAGTT ATGAGTG'TTT AATAATCAGT AATTACCAAT AAAGAATTAA	60
GAATACAAGT TTACAAGAAA TAAGTGTTAC AAAAAATAGC TGAAAAGGAA GATGAAC	117
ATG AAC AAG AAT AAT ACT AAA TTA AGC ACA AGA GCC TTA CCA AGT TTT	165
Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe	
790 795 800 805	

ATT GAT TAT TTC AAT GGC ATT TAT GGA TTT GCC ACT GGT ATC AAA GAC Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp 810 815 820	213
ATT ATG AAC ATG ATT TTT AAA ACG GAT ACA GGT GGT GAT CTA ACC CTA Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu 825 830 835	261
GAC GAA ATT TTA AAG AAT CAG CAG CTA CTA AAT GAT ATT TCT GGT AAA Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys 840 845 850	309
TTG GAT GGG GTG AAT GGA AGC TTA AAT GAT CTT ATC GCA CAG GGA AAC Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn 855 860 865	357
TTA AAT ACA GAA TTA TCT AAG GAA ATA TTA AAA ATT GCA AAT GAA CAA Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln 870 875 880 885	405
AAT CAA GTT TTA AAT GAT GTT AAT AAC AAA CTC GAT GCG ATA AAT ACG Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr 890 895 900	453
ATG CTT CGG GTA TAT CTA CCT AAA ATT ACC TCT ATG TTG AGT GAT GTA Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val 905 910 915	501
ATG AAA CAA AAT TAT GCG CTA AGT CTG CAA ATA GAA TAC TTA AGT AAA Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys 920 925 930	549
CAA TTG CAA GAG ATT TCT GAT AAG TTG GAT ATT ATT AAT GTA AAT GTA Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val 935 940 945	597
CTT ATT AAC TCT ACA CTT ACT GAA ATT ACA CCT GCG TAT CAA AGG ATT Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile 950 955 960 965	645
AAA TAT GTG AAC GAA AAA TTT GAG GAA TTA ACT TTT GCT ACA GAA ACT Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr 970 975 980	693
AGT TCA AAA GTA AAA AAG GAT GGC TCT CCT GCA GAT ATT CGT GAT GAG Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Arg Asp Glu 985 990 995	741
TTA ACT GAG TTA ACT GAA CTA GCG AAA AGT GTA ACA AAA AAT GAT GTG Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val 1000 1005 1010	789
GAT GGT TTT GAA TTT TAC CTT AAT ACA TTC CAC GAT GTA ATG GTA GGA Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly 1015 1020 1025	837
AAT AAT TTA TTC GGG CGT TCA GCT TTA AAA ACT GCA TCG GAA TTA ATT Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile 1030 1035 1040 1045	885
ACT AAA GAA AAT GTG AAA ACA AGT GGC AGT GAG GTC GGA AAT GTT TAT Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr 1050 1055 1060	933

AAC TTC CTA ATT GTA TTA ACA GCT CTG CAA GCA AAA GCT TTT CTT ACT Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr 1065 1070 1075	981
TTA ACA CCA TGC CGA AAA TTA TTA GGC TTA GCA GAT ATT GAT TAT ACT Leu Thr Pro Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr 1080 1085 1090	1029
TCT ATT ATG AAT GAA CAT TTA AAT AAG GAA AAA GAG GAA TTT AGA GTA Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val 1095 1100 1105	1077
AAC ATC CTC CCT ACA CTT TCT AAT ACT TTT TCT AAT CCT AAT TAT GCA Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala 1110 1115 1120 1125	1125
AAA GTT AAA GGA AGT GAT GAA GAT GCA AAG ATG ATT GTG GAA GCT AAA Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys 1130 1135 1140	1173
CCA GGA CAT GCA TTG ATT GGG TTT GAA ATT AGT AAT GAT TCA ATT ACA Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr 1145 1150 1155	1221
GTA TTA AAA GTA TAT GAG GCT AAG CTA AAA CAA AAT TAT CAA GTC GAT Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp 1160 1165 1170	1269
AAG GAT TCC TTA TCG GAA GTT ATT TAT GGC GAT ATG GAT AAA TTA TTG Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu 1175 1180 1185	1317
TGC CCA GAT CAA TCT GGA CAA ATC TAT TAT ACA AAT AAC ATA GTA TTT Cys Pro Asp Gln Ser Gly Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe 1190 1195 1200 1205	1365
CCA AAT GAA TAT GTA ATT ACT AAA ATT GAT TTC ACT AAA AAA ATG AAA Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys 1210 1215 1220	1413
ACT TTA AGA TAT GAG GTA ACA GCG AAT TTT TAT GAT TCT TCT ACA GGA Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly 1225 1230 1235	1461
GAA ATT GAC TTA AAT AAG AAA AAA GTA GAA TCA AGT GAA GCG GAG TAT Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr 1240 1245 1250	1509
AGA ACG TTA AGT GCT AAT GAT GAT GGG GTG TAT ATG CCG TTA GGT GTC Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val 1255 1260 1265	1557
ATC AGT GAA ACA TTT TTG ACT CCG ATT AAT GGG TTT GGC CTC CAA GCT Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala 1270 1275 1280 1285	1605
GAT GAA AAT TCA AGA TTA ATT ACT TTA ACA TGT AAA TCA TAT TTA AGA Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg 1290 1295 1300	1653
GAA CTA CTG CTA GCA ACA GAC TTA AGC AAT AAA GAA ACT AAA TTG ATC Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile 1305 1310 1315	1701
GTC CCG CCA AGT GGT TTT ATT AGC AAT ATT GTA GAG AAC GGG TCC ATA	1749



1575

TCTTGATATAA TGTCCTTGAT TATGGAAAAA CACAATTTTG TTTGCTAAGA TGTATATATA 2584

GCTCACTCAT TAAAAGGCAA TCAAGCTT 2612

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
 1              5              10              15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
          20              25              30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
          35              40              45

Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
          50              55              60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
          65              70              75              80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
          85              90              95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
          100              105              110

Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
          115              120              125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
          130              135              140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
          145              150              155              160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
          165              170              175

Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
          180              185              190

Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Arg Asp Glu
          195              200              205

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
          210              215              220

Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
          225              230              235              240

Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
          245              250              255

```



Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr  
 260 265 270  
 Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr  
 275 280 285  
 Leu Thr Pro Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr  
 290 295 300  
 Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val  
 305 310 315 320  
 Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala  
 325 330 335  
 Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys  
 340 345 350  
 Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr  
 355 360 365  
 Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp  
 370 375 380  
 Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu  
 385 390 395 400  
 Cys Pro Asp Gln Ser Gly Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe  
 405 410 415  
 Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys  
 420 425 430  
 Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly  
 435 440 445  
 Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr  
 450 455 460  
 Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val  
 465 470 475 480  
 Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala  
 485 490 495  
 Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg  
 500 505 510  
 Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile  
 515 520 525  
 Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile  
 530 535 540  
 Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr  
 545 550 555 560  
 Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His  
 565 570 575  
 Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys  
 580 585 590  
 Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His

- 68 -

595	600	605
Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn		
610	615	620
Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr		
625	630	635
Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu		
645	650	655
Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys		
660	665	670
Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly		
675	680	685
Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg		
690	695	700
Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg		
705	710	715
Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser		
725	730	735
Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val		
740	745	750
Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr Ile Glu		
755	760	765
Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His Phe Tyr		
770	775	780
Asp Val Ser Ile Lys		
785		

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2364 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 56..2295
- (D) OTHER INFORMATION: /product= "VIP3A(c) protein"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AATACAATTT ACGAGGGATA AGTGTACAA AGAATAGCTG AGGAGGGAGA TGAAC ATG  
Met  
1

58

AAC AAG AAT AAT GCT AAA TTA AGC ACA AGA GCC TTA CCA AGT TTT ATT Asn Lys Asn Asn Ala Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe Ile	106
5 10 15	
GAT TAT TTC AAT GGC ATT TAT GGA TTT GCC ACT GGT ATC AAA GAC ATT Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp Ile	154
20 25 30	
ATG AAC ATG ATT TTT AAA ACG GAT ACA GGT GGT GAT CTA GCC CTA GAC Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Ala Leu Asp	202
35 40 45	
GAA ATT TTA GAG AAT CAG CAG CTA CTA AAT GAT ATT TCT GGT AAA TTG Glu Ile Leu Glu Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys Leu	250
50 55 60 65	
GAT GGG GTG AAT GGA AGC TTA AAT GAT CTT ATC GCA CAG GGA AAC TTA Asp Gly Val Asn Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn Leu	298
70 75 80	
AAT ACA GAA TTA TCT AAG GAA ATA TTA AAA ATT GCA AAT GAA CAA AAT Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln Asn	346
85 90 95	
CAA GTT TTA AAT GAT GTT AAT AAC AAA CTC GAT GCG ATA AAT ACG ATG Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr Met	394
100 105 110	
CTT CGG GTA TAT CTA CCT AAA ATT ACC TCT ATG TTG AGT GAT GTA ATG Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val Met	442
115 120 125	
AAA CAA AAT TAT GCG CTA AGT CTG CAA ATA GAA TAC TTA AGT AAA CAA Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys Gln	490
130 135 140 145	
TTG CAA GAG ATT TCT GAT AAG TTG GAT ATT ATT AAT GTA AAT GTA CTT Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val Leu	538
150 155 160	
ATT AAC TCT ACA CTT ACT GAA ATT ACA CCT GCG TAT CAA AGG ATT AAA Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile Lys	586
165 170 175	
TAT GTG AAC GAA AAA TTT GAG GAA TTA ACT TTT GCT ACA GAA ACT AGT Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr Ser	634
180 185 190	
TCA AAA GTA AAA AAG GAT GGC TCT CCT GCA GAT ATT CGT GAT GAG TTA Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Arg Asp Glu Leu	682
195 200 205	
AGT GAG TTA ACT GAA CTA GCG AAA AGT GTA ACA CAA AAT GAT GTG GAT Ser Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Gln Asn Asp Val Asp	730
210 215 220 225	
GGT TTT GAA TTT TAC CTT AAT ACA TTC CAC GAT GTA ATG GTA GGA AAT Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly Asn	778
230 235 240	

AAT TTA TTC GGG CGT TCA GCT TTA AAA ACT GCA TCG GAA TTA ATT ACT	826
Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile Thr	
245 250 255	
AAA GAA AAT GTG AAA ACA AGT GGC AGT GAG GTC GGA AAT GTT TAT AAC	874
Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr Asn	
260 265 270	
TTC CTA ATT GTA TTA ACA GCT CTG CAA GCA CAA GCT TTT CTT ACT TTA	922
Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Gln Ala Phe Leu Thr Leu	
275 280 285	
ACA CCA TGC CGA AAA TTA TTA GGC TTA GCA GAT ATT GAT TAT ACT TCT	970
Thr Pro Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr Ser	
290 295 300 305	
ATT ATG AAT GAA CAT TTA AAT AAG GAA AAA GAG GAA TTT AGA GTA AAC	1018
Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val Asn	
310 315 320	
ATC CTC CCT ACA CTT TCT AAT ACT TTT TCT AAT CCT AAT TAT GCA AAA	1066
Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala Lys	
325 330 335	
GTT AAA GGA AGT GAT GAA GAT GCA AAG ATG ATT GTG GAA GCT AAA CCA	1114
Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys Pro	
340 345 350	
GGA CAT GCA TTG ATT GGG TTT GAA ATT AGT AAT GAT TCA ATT ACA GTA	1162
Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr Val	
355 360 365	
TTA AAA GTA TAT GAG GCT AAG CTA AAA CAA AAT TAT CAA GTC GAT AAG	1210
Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp Lys	
370 375 380 385	
GAT TCC TTA TCG GAA GTT ATT TAT GGC GAT ATG GAT AAA TTA TTG TGC	1258
Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu Cys	
390 395 400	
CCA GAT CAA TCT GGA CAA ATC TAT TAT ACA AAT AAC ATA GTA TTT CCA	1306
Pro Asp Gln Ser Gly Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe Pro	
405 410 415	
AAT GAA TAT GTA ATT ACT AAA ATT GAT TTC ACT AAA AAA ATG AAA ACT	1354
Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys Thr	
420 425 430	
TTA AGA TAT GAG GTA ACA GCG AAT TTT TAT GAT TCT TCT ACA GGA GAA	1402
Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly Glu	
435 440 445	
ATT GAC TTA AAT AAG AAA AAA GTA GAA TCA AGT GAA GCG GAG TAT AGA	1450
Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr Arg	
450 455 460 465	
ACG TTA AGT GCT AAT GAT GAT GGG GTG TAT ATG CCG TTA GGT GTC ATC	1498
Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val Ile	
470 475 480	

AGT GAA ACA TTT TTG ACT CCG ATT AAT GGG TTT GGC CTC CAA GCT GAT Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala Asp 485 490 495	1546
GAA AAT TCA AGA TTA ATT ACT TTA ACA TGT AAA TCA TAT TTA AGA GAA Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg Glu 500 505 510	1594
CTA CTG CTA GCA ACA GAC TTA AGC AAT AAA GAA ACT AAA TTG ATC GTC Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile Val 515 520 525	1642
CCG CCA AGT GGT TTT ATT AGC AAT ATT GTA GAG AAC GGG TCC ATA GAA Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile Glu 530 535 540 545	1690
GAG GAC AAT TTA GAG CCG TGG AAA GCA AAT AAT AAG AAT GCG TAT GTA Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Lys Asn Ala Tyr Val 550 555 560	1738
GAT CAT ACA GGC GGA GTG AAT GGA ACT AAA GCT TTA TAT GTT CAT AAG Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His Lys 565 570 575	1786
GAC GGA GGA ATT TCA CAA TTT ATT GGA GAT AAG TTA AAA CCG AAA ACT Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys Thr 580 585 590	1834
GAG TAT GTA ATC CAA TAT ACT GTT AAA GGA AAA CCT TCT ATT CAT TTA Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His Leu 595 600 605	1882
AAA GAT GAA AAT ACT GGA TAT ATT CAT TAT GAA GAT ACA AAT AAT AAT Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn Asn 610 615 620 625	1930
TTA GAA GAT TAT CAA ACT ATT AAT AAA CGT TTT ACT ACA GGA ACT GAT Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr Asp 630 635 640	1978
TTA AAG GGA GTG TAT TTA ATT TTA AAA AGT CAA AAT GGA GAT GAA GCT Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu Ala 645 650 655	2026
TGG GGA GAT AAC TTT ATT ATT TTG GAA ATT AGT CCT TCT GAA AAG TTA Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys Leu 660 665 670	2074
TTA AGT CCA GAA TTA ATT AAT ACA AAT AAT TGG ACG AGT ACG GGA TCA Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly Ser 675 680 685	2122
ACT AAT ATT AGC GGT AAT ACA CTC ACT CTT TAT CAG GGA GGA CGA GGG Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg Gly 690 695 700 705	2170
ATT CTA AAA CAA AAC CTT CAA TTA GAT AGT TTT TCA ACT TAT AGA GTG Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg Val 710 715 720	2218

TAT TTC TCT GTG TCC GGA GAT GCT AAT GTA AGG ATT AGA AAT TCT AGG 2266  
 Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser Arg  
           725                              730                              735

GAA GTG TTA TTT GAA AAA AAG GAT ATA TGA GC GCGCTAAAG ATGTTTCTGA 2318  
 Glu Val Leu Phe Glu Lys Lys Asp Ile  
           740                              745

AATGTTCACT ACAAATTGA AAGATAACTT CTATATAGAG CTTTCT 2364

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 747 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asn Lys Asn Asn Ala Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe  
   1                  5                  10                  15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp  
           20                  25                  30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Ala Leu  
           35                  40                  45

Asp Glu Ile Leu Glu Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys  
           50                  55                  60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn  
           65                  70                  75                  80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln  
           85                  90                  95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr  
           100                  105                  110

Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val  
           115                  120                  125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys  
           130                  135                  140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val  
           145                  150                  155                  160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile  
           165                  170                  175

Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr  
           180                  185                  190

Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Arg Asp Glu

- 73 -

195					200					205					
Leu	Ser	Glu	Leu	Thr	Glu	Leu	Ala	Lys	Ser	Val	Thr	Gln	Asn	Asp	Val
210					215					220					
Asp	Gly	Phe	Glu	Phe	Tyr	Leu	Asn	Thr	Phe	His	Asp	Val	Met	Val	Gly
225					230					235					240
Asn	Asn	Leu	Phe	Gly	Arg	Ser	Ala	Leu	Lys	Thr	Ala	Ser	Glu	Leu	Ile
				245					250					255	
Thr	Lys	Glu	Asn	Val	Lys	Thr	Ser	Gly	Ser	Glu	Val	Gly	Asn	Val	Tyr
			260					265					270		
Asn	Phe	Leu	Ile	Val	Leu	Thr	Ala	Leu	Gln	Ala	Gln	Ala	Phe	Leu	Thr
		275					280					285			
Leu	Thr	Pro	Cys	Arg	Lys	Leu	Leu	Gly	Leu	Ala	Asp	Ile	Asp	Tyr	Thr
		290				295					300				
Ser	Ile	Met	Asn	Glu	His	Leu	Asn	Lys	Glu	Lys	Glu	Glu	Phe	Arg	Val
305					310					315					320
Asn	Ile	Leu	Pro	Thr	Leu	Ser	Asn	Thr	Phe	Ser	Asn	Pro	Asn	Tyr	Ala
				325					330					335	
Lys	Val	Lys	Gly	Ser	Asp	Glu	Asp	Ala	Lys	Met	Ile	Val	Glu	Ala	Lys
			340					345					350		
Pro	Gly	His	Ala	Leu	Ile	Gly	Phe	Glu	Ile	Ser	Asn	Asp	Ser	Ile	Thr
		355				360						365			
Val	Leu	Lys	Val	Tyr	Glu	Ala	Lys	Leu	Lys	Gln	Asn	Tyr	Gln	Val	Asp
	370					375					380				
Lys	Asp	Ser	Leu	Ser	Glu	Val	Ile	Tyr	Gly	Asp	Met	Asp	Lys	Leu	Leu
385					390					395					400
Cys	Pro	Asp	Gln	Ser	Gly	Gln	Ile	Tyr	Tyr	Thr	Asn	Asn	Ile	Val	Phe
			405						410					415	
Pro	Asn	Glu	Tyr	Val	Ile	Thr	Lys	Ile	Asp	Phe	Thr	Lys	Lys	Met	Lys
			420				425						430		
Thr	Leu	Arg	Tyr	Glu	Val	Thr	Ala	Asn	Phe	Tyr	Asp	Ser	Ser	Thr	Gly
		435					440					445			
Glu	Ile	Asp	Leu	Asn	Lys	Lys	Lys	Val	Glu	Ser	Ser	Glu	Ala	Glu	Tyr
	450					455					460				
Arg	Thr	Leu	Ser	Ala	Asn	Asp	Asp	Gly	Val	Tyr	Met	Pro	Leu	Gly	Val
465					470					475					480
Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Asn	Gly	Phe	Gly	Leu	Gln	Ala
			485						490					495	
Asp	Glu	Asn	Ser	Arg	Leu	Ile	Thr	Leu	Thr	Cys	Lys	Ser	Tyr	Leu	Arg
			500					505					510		
Glu	Leu	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Glu	Thr	Lys	Leu	Ile

515					520					525					
Val	Pro	Pro	Ser	Gly	Phe	Ile	Ser	Asn	Ile	Val	Glu	Asn	Gly	Ser	Ile
530						535					540				
Glu	Glu	Asp	Asn	Leu	Glu	Pro	Trp	Lys	Ala	Asn	Asn	Lys	Asn	Ala	Tyr
545					550					555					560
Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Ala	Leu	Tyr	Val	His
				565					570					575	
Lys	Asp	Gly	Gly	Ile	Ser	Gln	Phe	Ile	Gly	Asp	Lys	Leu	Lys	Pro	Lys
				580					585					590	
Thr	Glu	Tyr	Val	Ile	Gln	Tyr	Thr	Val	Lys	Gly	Lys	Pro	Ser	Ile	His
		595					600					605			
Leu	Lys	Asp	Glu	Asn	Thr	Gly	Tyr	Ile	His	Tyr	Glu	Asp	Thr	Asn	Asn
610						615					620				
Asn	Leu	Glu	Asp	Tyr	Gln	Thr	Ile	Asn	Lys	Arg	Phe	Thr	Thr	Gly	Thr
625					630					635					640
Asp	Leu	Lys	Gly	Val	Tyr	Leu	Ile	Leu	Lys	Ser	Gln	Asn	Gly	Asp	Glu
				645					650					655	
Ala	Trp	Gly	Asp	Asn	Phe	Ile	Ile	Leu	Glu	Ile	Ser	Pro	Ser	Glu	Lys
				660				665						670	
Leu	Leu	Ser	Pro	Glu	Leu	Ile	Asn	Thr	Asn	Asn	Trp	Thr	Ser	Thr	Gly
		675					680					685			
Ser	Thr	Asn	Ile	Ser	Gly	Asn	Thr	Leu	Thr	Leu	Tyr	Gln	Gly	Gly	Arg
		690				695					700				
Gly	Ile	Leu	Lys	Gln	Asn	Leu	Gln	Leu	Asp	Ser	Phe	Ser	Thr	Tyr	Arg
705				710						715					720
Val	Tyr	Phe	Ser	Val	Ser	Gly	Asp	Ala	Asn	Val	Arg	Ile	Arg	Asn	Ser
				725					730					735	
Arg	Glu	Val	Leu	Phe	Glu	Lys	Lys	Asp	Ile						
			740					745							

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2403 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Synthetic DNA"

## (iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 11..2389



(D) OTHER INFORMATION: /note= "maize optimized DNA  
sequence encoding VIP3A(a) "

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGATCCACCA ATGAACATGA ACAAGAACAA CACCAAGCTG AGCACCCGCG CCCTGCCGAG	60
CTTCATCGAC TACTTCAACG GCATCTACGG CTTGCCACC GGCATCAAGG ACATCATGAA	120
CATGATCTTC AAGACCGACA CCGGCGGCGA CCTGACCCTG GACGAGATCC TGAAGAACCA	180
GCAGCTGCTG AACGACATCA GCGGCAAGCT GGACGGCGTG AACGGCAGCC TGAACGACCT	240
GATCGCCCAG GGCAACCTGA ACACCGAGCT GAGCAAGGAG ATCCTTAAGA TCGCCAACGA	300
GCAGAACCAG GTGCTGAACG ACGTGAACAA CAAGCTGGAC GCCATCAACA CCATGCTGCG	360
CGTGTACCTG CCGAAGATCA CCAGCATGCT GAGCGACGTG ATGAAGCAGA ACTACGCCCT	420
GAGCCTGCAG ATCGAGTACC TGAGCAAGCA GCTGCAGGAG ATCAGCGACA AGCTGGACAT	480
CATCAACGTG AACGTCTCTGA TCAACAGCAC CCTGACCGAG ATCACCCCGG CCTACCAGCG	540
CATCAAGTAC GTGAACGAGA AGTTCGAAGA GCTGACCTTC GCCACCGAGA CCAGCAGCAA	600
GGTGAAGAAG GACGGCAGCC CGGCCGACAT CCTGGACGAG CTGACCGAGC TGACCGAGCT	660
GGCCAAGAGC GTGACCAAGA ACGACGTGGA CGGCTTCGAG TTCTACCTGA ACACCTTCCA	720
CGACGTGATG GTGGGCAACA ACCTGTTCGG CCGCAGCGCC CTGAAGACCG CCAGCGAGCT	780
GATCACCAAG GAGAACGTGA AGACCAGCGG CAGCGAGGTG GGCAACGTGT ACAACTTCCT	840
GATCGTGCTG ACCGCCCTGC AGGCCAGGC CTTCTGACC CTGACCACCT GTCGCAAGCT	900
GCTGGGCCTG GCCGACATCG ACTACACCAG CATCATGAAC GAGCACTTGA ACAAGGAGAA	960
GGAGGAGTTC CGCGTGAACA TCCTGCCGAC CCTGAGCAAC ACCTTCAGCA ACCCGAACTA	1020
CGCCAAGGTG AAGGGCAGCG ACGAGGACGC CAAGATGATC GTGGAGGCTA AGCCGGGCCA	1080
CGCGTTGATC GGCTTCGAGA TCAGCAACGA CAGCATCACC GTGCTGAAGG TGTACGAGGC	1140
CAAGCTGAAG CAGAACTACC AGGTGGACAA GGACAGCTTG AGCGAGGTGA TCTACGGCGA	1200
CATGGACAAG CTGCTGTGTC CGGACCAGAG CGAGCAAATC TACTACACCA ACAACATCGT	1260
GTTCCCGAAC GAGTACGTGA TCACCAAGAT CGACTTCACC AAGAAGATGA AGACCCTGCG	1320
CTACGAGGTG ACCGCCAACT TCTACGACAG CAGCACCAGC GAGATCGACC TGAACAAGAA	1380
GAAGGTGGAG AGCAGCGAGG CCGAGTACCG CACCCTGAGC GCGAACGACG ACGGCGTCTA	1440
CATGCCACTG GCGGTGATCA GCGAGACCTT CCTGACCCCG ATCAACGGCT TTGGCCTGCA	1500
GGCCGACGAG AACAGCCGCC TGATCACCCT GACCTGTAAG AGCTACCTGC GCGAGCTGCT	1560
GCTAGCCACC GACCTGAGCA ACAAGGAGAC CAAGCTGATC GTGCCACCGA GCGGCTTCAT	1620
CAGCAACATC GTGGAGAACG GCAGCATCGA GGAGGACAAC CTGGAGCCGT GGAAGGCCAA	1680
CAACAAGAAC GCCTACGTGG ACCACACCGG CGGCGTGAAC GGCACCAAGG CCCTGTACGT	1740
GCACAAGGAC GGCGGCATCA GCCAGTTCAT CGGCGACAAG CTGAAGCCGA AGACCGAGTA	1800

```

CGTGATCCAG TACACCGTGA AGGGCAAGCC ATCGATTAC CTGAAGGACG AGAACACCGG      1860
CTACATCCAC TACGAGGACA CCAACAACAA CCTGGAGGAC TACCAGACCA TCAACAAGCG      1920
CTTCACCACC GGCACCGACC TGAAGGGCGT GTACCTGATC CTGAAGAGCC AGAACGGCGA      1980
CGAGGCCTGG GCGGACAAC TCAATCATCCT GGAGATCAGC CCGAGCGAGA AGCTGCTGAG      2040
CCCGGAGCTG ATCAACACCA ACAACTGGAC CAGCACCGGC AGCACCAACA TCAGCGGCAA      2100
CACCTGACC CTGTACCAGG GCGGCCGCGG CATCCTGAAG CAGAACCTGC AGCTGGACAG      2160
CTTCAGCACC TACCGCGTGT ACTTCAGCGT GAGCGGCGAC GCCAACCTGC GCATCCGCAA      2220
CAGCCGCGAG GTGCTGTTTC AGAAGAGGTA CATGAGCGGC GCCAAGGACG TGAGCGAGAT      2280
GTTACCACC AAGTTCGAGA AGGACAACCT CTACATCGAG CTGAGCCAGG GCAACAACCT      2340
GTACGGCGGC CCGATCGTGC ACTTCTACGA CGTGAGCATC AAGTTAACGT AGAGCTCAGA      2400
TCT                                                                                   2403

```

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1638 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 44..1191
- (D) OTHER INFORMATION: /product= "Translation of cDNA encoding VIP3A(a) receptor"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

T AGT GGA TCC CCC GGG CTG CAG GAA TTC GCG GCC GCG TCG ACC ATG      46
                               Met
                               15
1           5           10
TAC TCT AGA ATA TTT TTC CTC CTT GTG ATA GTG TGT GCT GTT AAG GCT      94
Tyr Ser Arg Ile Phe Phe Leu Leu Val Ile Val Cys Ala Val Lys Ala
                20                25                30
TCT CTG TTT ACT GTA AAT GTG TAT GAT GAT AAC CCC GAA ACT GAA ATT      142
Ser Leu Phe Thr Val Asn Val Tyr Asp Asp Asn Pro Glu Thr Glu Ile
                35                40                45
GCG AGT AGT CTA AAA GGC TGT AAC CCC CAA GAG TGT GAC CAG CGG TGT      190
Ala Ser Ser Leu Lys Gly Cys Asn Pro Gln Glu Cys Asp Gln Arg Cys
                50                55                60

```

CGT AGA CTG AAG TTT CCC GGT GGC GCC TGT GTC AAT GGT CGC TGC AAG Arg Arg Leu Lys Phe Pro Gly Gly Ala Cys Val Asn Gly Arg Cys Lys 65 70 75	238
TGT GAC AAC TTC CTC AGT GTA AAA GAT GAC GTG TCT GTT GAA GAG CCT Cys Asp Asn Phe Leu Ser Val Lys Asp Asp Val Ser Val Glu Glu Pro 80 85 90 95	286
GCG ATT CTC AAA GAT TTG GTG TCA TTA GAA GCT GAA CAG GCA GCG AAA Ala Ile Leu Lys Asp Leu Val Ser Leu Glu Ala Glu Gln Ala Ala Lys 100 105 110	334
AGT AGA TGC AGA AAC AGA GTG TGT GAC GCG GTG TGC CGT GCC CTA CAC Ser Arg Cys Arg Asn Arg Val Cys Asp Ala Val Cys Arg Ala Leu His 115 120 125	382
AAC ACC AGT GGT GCC TGT GTT GAT GGA CAA TGC AAG TGT ACT AAT AAG Asn Thr Ser Gly Ala Cys Val Asp Gly Gln Cys Lys Cys Thr Asn Lys 130 135 140	430
ATC AGT GCA GGA GAT ATT GTG TCT GAT CCT GCT GAA TCG CTA CGC ACT Ile Ser Ala Gly Asp Ile Val Ser Asp Pro Ala Glu Ser Leu Arg Thr 145 150 155	478
TGT AAC CCT ATA AGG TGT GAC GAA CAA TGT AGA AGA AAT GGC CAT GAA Cys Asn Pro Ile Arg Cys Asp Glu Gln Cys Arg Arg Asn Gly His Glu 160 165 170 175	526
TTT GGT GTT TGC TTC AAA GGA CAA TGC AAG TGT GAT TAC TTC CTC AAG Phe Gly Val Cys Phe Lys Gly Gln Cys Lys Cys Asp Tyr Phe Leu Lys 180 185 190	574
GAA GAA GTC GAT GAA CCT GAA GTT ACA AGC CTT CCA AAA AAC TGC AAC Glu Glu Val Asp Glu Pro Glu Val Thr Ser Leu Pro Lys Asn Cys Asn 195 200 205	622
CCC CAA GAG TGT GAC CAG CGT TGT CGT AGA CTG AAG TTC CCC GGT GGC Pro Gln Glu Cys Asp Gln Arg Cys Arg Arg Leu Lys Phe Pro Gly Gly 210 215 220	670
GCC TGT GTC AAC GGG CGC TGC AAG TGT GAC AAC TTC TTC AGT GCA GGA Ala Cys Val Asn Gly Arg Cys Lys Cys Asp Asn Phe Phe Ser Ala Gly 225 230 235	718
GAT ATT GTG TCT GAT CCT GCC GAA TCG CTA CGC TCT TGT AAC CCT ATA Asp Ile Val Ser Asp Pro Ala Glu Ser Leu Arg Ser Cys Asn Pro Ile 240 245 250 255	766
AGG TGT GAC GAA CAA TGT AGA AGA AAT GGC CAT GAA TTT GGT GTT TGC Arg Cys Asp Glu Gln Cys Arg Arg Asn Gly His Glu Phe Gly Val Cys 260 265 270	814
TTC AAA GGA CAA TGC AAG TGT GAT TAC TTC CTC AAC TCA GAA GTA GAC Phe Lys Gly Gln Cys Lys Cys Asp Tyr Phe Leu Asn Ser Glu Val Asp 275 280 285	862
GCT GTT AAT GAG TTT CCT CAA GCG GGC TCA AAA CGC TAC TGC AAC TTA Ala Val Asn Glu Phe Pro Gln Ala Gly Ser Lys Arg Tyr Cys Asn Leu 290 295 300	910

ACG CAA TGC AAC CAG ACG TGC GCC AAT CGT TTC TAT GAT AGT GCT AGA 958  
 Thr Gln Cys Asn Gln Thr Cys Ala Asn Arg Phe Tyr Asp Ser Ala Arg  
 305 310 315  
 GTG ATC CAC GGC TGG TGC AAA TGC TAC AGT AAG ATG GAA AGA CAG GAT 1006  
 Val Ile His Gly Trp Cys Lys Cys Tyr Ser Lys Met Glu Arg Gln Asp  
 320 325 330 335  
 GCA TCT CCA TTA AAC GAT GTG ACT GAG GAT GAA AAT GAA GTT TCT AAC 1054  
 Ala Ser Pro Leu Asn Asp Val Thr Glu Asp Glu Asn Glu Val Ser Asn  
 340 345 350  
 GAT ATC CTG AGG ACT GTT GCA GAG GAG CTG TCT GAT GTG TCA CCT AGG 1102  
 Asp Ile Leu Arg Thr Val Ala Glu Glu Leu Ser Asp Val Ser Pro Arg  
 355 360 365  
 GCC TGC AAA TCA GCG AGC TGC AAT CAA GCA TGT CGC GCC TTC TAC TTT 1150  
 Ala Cys Lys Ser Ala Ser Cys Asn Gln Ala Cys Arg Ala Phe Tyr Phe  
 370 375 380  
 AAA GGA GGG TGG TGT CGC TTT GGA CGA TGC CAA TGC TTC TA 1191  
 Lys Gly Gly Trp Cys Arg Phe Gly Arg Cys Gln Cys Phe  
 385 390 395  
 AAATTAGTAT GATATATGAA TTTTGTATTA TTCGGTTAAT TGTGTTATGT TTAAAAAACA 1251  
 TAATGTCTTC ATTTTAGAAA AAAGTACCTT CACTAAAGCG CAACAATTAA CTAGTAGTTA 1311  
 ATTATTAACT AGTAGTTAAA TTATTGATGA TTATGATTAT CTTAGTAGTA GTTAATTATA 1371  
 ATCATCAACT ATTAAC TAGT AGTTAATTAT TAAC TAGTAG TTAAATTATT GATGATTATG 1431  
 ATTATCTTAG TAGTAGTTAA TTATTGTTTC TTATAATAAT CTAGTATGTT GGTAGGTACT 1491  
 TAATAATAAC GCTTCTGACA AAAAATTTAA AATTAAATAA TTCTATCAAA CATAAATAAT 1551  
 AACTGAAATA AAAATTTATA AGAGAAAAAA AAAAAGTCGA CGCGGCCGCG AATTCGATAT 1611  
 CAAGCTTATC GATACCGTCG ACCTCGA 1638

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ser Gly Ser Pro Gly Leu Gln Glu Phe Ala Ala Ala Ser Thr Met Tyr  
 1 5 10 15  
 Ser Arg Ile Phe Phe Leu Leu Val Ile Val Cys Ala Val Lys Ala Ser  
 20 25 30  
 Leu Phe Thr Val Asn Val Tyr Asp Asp Asn Pro Glu Thr Glu Ile Ala  
 35 40 45

Ser Ser Leu Lys Gly Cys Asn Pro Gln Glu Cys Asp Gln Arg Cys Arg  
 50 55 60  
 Arg Leu Lys Phe Pro Gly Gly Ala Cys Val Asn Gly Arg Cys Lys Cys  
 65 70 75 80  
 Asp Asn Phe Leu Ser Val Lys Asp Asp Val Ser Val Glu Glu Pro Ala  
 85 90 95  
 Ile Leu Lys Asp Leu Val Ser Leu Glu Ala Glu Gln Ala Ala Lys Ser  
 100 105 110  
 Arg Cys Arg Asn Arg Val Cys Asp Ala Val Cys Arg Ala Leu His Asn  
 115 120 125  
 Thr Ser Gly Ala Cys Val Asp Gly Gln Cys Lys Cys Thr Asn Lys Ile  
 130 135 140  
 Ser Ala Gly Asp Ile Val Ser Asp Pro Ala Glu Ser Leu Arg Thr Cys  
 145 150 155 160  
 Asn Pro Ile Arg Cys Asp Glu Gln Cys Arg Arg Asn Gly His Glu Phe  
 165 170 175  
 Gly Val Cys Phe Lys Gly Gln Cys Lys Cys Asp Tyr Phe Leu Lys Glu  
 180 185 190  
 Glu Val Asp Glu Pro Glu Val Thr Ser Leu Pro Lys Asn Cys Asn Pro  
 195 200 205  
 Gln Glu Cys Asp Gln Arg Cys Arg Arg Leu Lys Phe Pro Gly Gly Ala  
 210 215 220  
 Cys Val Asn Gly Arg Cys Lys Cys Asp Asn Phe Phe Ser Ala Gly Asp  
 225 230 235 240  
 Ile Val Ser Asp Pro Ala Glu Ser Leu Arg Ser Cys Asn Pro Ile Arg  
 245 250 255  
 Cys Asp Glu Gln Cys Arg Arg Asn Gly His Glu Phe Gly Val Cys Phe  
 260 265 270  
 Lys Gly Gln Cys Lys Cys Asp Tyr Phe Leu Asn Ser Glu Val Asp Ala  
 275 280 285  
 Val Asn Glu Phe Pro Gln Ala Gly Ser Lys Arg Tyr Cys Asn Leu Thr  
 290 295 300  
 Gln Cys Asn Gln Thr Cys Ala Asn Arg Phe Tyr Asp Ser Ala Arg Val  
 305 310 315 320  
 Ile His Gly Trp Cys Lys Cys Tyr Ser Lys Met Glu Arg Gln Asp Ala  
 325 330 335  
 Ser Pro Leu Asn Asp Val Thr Glu Asp Glu Asn Glu Val Ser Asn Asp  
 340 345 350  
 Ile Leu Arg Thr Val Ala Glu Glu Leu Ser Asp Val Ser Pro Arg Ala  
 355 360 365

- 80 -

Cys Lys Ser Ala Ser Cys Asn Gln Ala Cys Arg Ala Phe Tyr Phe Lys  
370 375 380

Gly Gly Trp Cys Arg Phe Gly Arg Cys Gln Cys Phe  
385 390 395

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

- (vi) ORIGINAL SOURCE:  
(A) ORGANISM: *Bacillus thuringiensis*  
(B) STRAIN: AB88

- (ix) FEATURE:  
(A) NAME/KEY: Peptide  
(B) LOCATION: 1..14  
(D) OTHER INFORMATION: /note= "N-terminal amino acid  
sequence of protein known as anion exchange fraction 23  
(smaller)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Xaa Glu Pro Phe Val Ser Ala Xaa Xaa Xaa Gln Xaa Xaa Xaa  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (vi) ORIGINAL SOURCE:  
(A) ORGANISM: *Bacillus thuringiensis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Xaa Glu Tyr Glu Asn Val Glu Pro Phe Val Ser Ala Xaa  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Bacillus thurigiensis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met	Asn	Lys	Asn	Asn	Thr	Lys	Leu	Pro	Thr	Arg	Ala	Leu	Pro
1				5					10				

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Bacillus thuringiensis*

(B) STRAIN: AB88

(ix) FEATURE:

(A) NAME/KEY: Peptide

(B) LOCATION: 1..15

(D) OTHER INFORMATION: /note= "N-terminal amino acid  
sequence of 35 kDa VIP active against *Agrotis ipsilon*"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ala	Leu	Ser	Glu	Asn	Thr	Gly	Lys	Asp	Gly	Gly	Tyr	Ile	Val	Pro
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Bacillus thuringiensis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met	Asp	Asn	Asn	Pro	Asn	Ile	Asn	Glu
1				5				

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..9
- (D) OTHER INFORMATION: /note= "N-terminal sequence of 80 kDa delta-endotoxin"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Met Asp Asn Asn Pro Asn Ile Asn Glu  
1 5

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 11 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Bacillus thuringiensis*

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /note= "N-terminal sequence from 60 kDa delta-endotoxin"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Met Asn Val Leu Asn Ser Gly Arg Thr Thr Ile  
1 5 10

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer sequence"



(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGATTAATGT TGGCCTC

17

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer sequence"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CATTAGCATC TCCGGACACA G

21

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2370 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA encoding VIP3A(b)"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ATGAACAAGA ACAACACCAA GCTGAGCACC CGCGCCCTGC CGAGCTTCAT CGACTACTTC	60
AACGGCATCT ACGGCTTCGC CACCGGCATC AAGGACATCA TGAACATGAT CTTCAAGACC	120
GACACCGGCG GCGACCTGAC CCTGGACGAG ATCCTGAAGA ACCAGCAGCT GCTGAACGAC	180
ATCAGCGGCA AGCTGGACGG CGTGAACGGC AGCCTGAACG ACCTGATCGC CCAGGGCAAC	240
CTGAACACCG AGCTGAGCAA GGAGATCCTT AAGATCGCCA ACGAGCAGAA CCAGGTGCTG	300
AACGACGTGA ACAACAAGCT GGACGCCATC AACACCATGC TGC GCGTGTA CCTGCCGAAG	360

ATCACCAGCA TGCTGAGCGA CGTGATGAAG CAGAACTACG CCCTGAGCCT GCAGATCGAG	420
TACCTGAGCA AGCAGCTGCA GGAGATCAGC GACAAGCTGG ACATCATCAA CGTGAACGTC	480
CTGATCAACA GCACCCTGAC CGAGATCACC CCGGCCTACC AGCGCATCAA GTACGTGAAC	540
GAGAAGTTCG AAGAGCTGAC CTTCGCCACC GAGACCAGCA GCAAGGTGAA GAAGGACGGC	600
AGCCCGGCCG ACATCCTGGA CGAGCTGACC GAGCTGACCG AGCTGGCGAA GAGCGTGACC	660
AAGAACGACG TGGACGGCTT CGAGTTCTAC CTGAACACCT TCCACGACGT GATGGTGGGC	720
AACAACCTGT TCGGCCGAG CGCCCTGAAG ACCGCCAGCG AGCTGATCAC CAAGGAGAAC	780
GTGAAGACCA GCGGCAGCGA GGTGGGCAAC GTGTACAACT TCCTGATCGT GCTGACCGCC	840
CTGCAGGCCA AGGCCTTCCT GACCCTGACC CCCTGTCGCA AGCTGCTGGG CCTGGCCGAC	900
ATCGACTACA CCAGCATCAT GAACGAGCAC TTGAACAAGG AGAAGGAGGA GTTCCGCGTG	960
AACATCCTGC CGACCCTGAG CAACACCTTC AGCAACCCGA ACTACGCCAA GGTGAAGGGC	1020
AGCGACGAGG ACGCCAAGAT GATCGTGGAG GCTAAGCCGG GCCACGCGTT GATCGGCTTC	1080
GAGATCAGCA ACGACAGCAT CACCGTGCTG AAGGTGTACG AGGCCAAGCT GAAGCAGAAC	1140
TACCAGGTGG ACAAGGACAG CTTGAGCGAG GTGATCTACG GCGACATGGA CAAGCTGCTG	1200
TGTCCGGACC AGAGCGGGCA AATCTACTAC ACCAACAACA TCGTGTTCCT GAACGAGTAC	1260
GTGATCACCA AGATCGACTT CACCAAGAAG ATGAAGACCC TCGCTACGA GGTGACCGCC	1320
AACTTCTACG ACAGCAGCAC CGGCGAGATC GACCTGAACA AGAAGAAGGT GGAGAGCAGC	1380
GAGGCCGAGT ACCGCACCCT GAGCGCGAAC GACGACGGCG TCTACATGCC ACTGGGCGTG	1440
ATCAGCGAGA CCTTCCTGAC CCCGATCAAC GGCTTTGGCC TGCAGGCCGA CGAGAACAGC	1500
CGCCTGATCA CCCTGACCTG TAAGAGCTAC CTGCGCGAGC TGCTGCTAGC CACCGACCTG	1560
AGCAACAAGG AGACCAAGCT GATCGTGCCA CCGAGCGGCT TCATCAGCAA CATCGTGGAG	1620
AACGGCAGCA TCGAGGAGGA CAACCTGGAG CCGTGGAAGG CCAACAACAA GAACGCCTAC	1680
GTGGACCACA CCGCGGGCGT GAACGGCACC AAGGCCCTGT ACGTGACAA GGACGGCGGC	1740
ATCAGCCAGT TCATCGGCGA CAAGCTGAAG CCGAAGACCG AGTACGTGAT CCAGTACACC	1800
GTGAAGGGCA AGCCATCGAT TCACCTGAAG GACGAGAACA CCGGCTACAT CCACTACGAG	1860
GACACCAACA ACAACCTGGA GGACTIONAG ACCATCAACA AGCGCTTCAC CACCGGCACC	1920
GACCTGAAGG GCGTGTACCT GATCCTGAAG AGCCAGAACG GCGACGAGGC CTGGGGCGAC	1980
AACTTCATCA TCCTGGAGAT CAGCCCGAGC GAGAAGCTGC TGAGCCCGGA GCTGATCAAC	2040
ACCAACAACCT GGACCAGCAC CGGCAGCACC AACATCAGCG GCAACACCCT GACCCTGTAC	2100
CAGGGCGGCC GCGGCATCCT GAAGCAGAAC CTGCAGCTGG ACAGCTTCAG CACCTACCGC	2160

GTGTACTTCA GCGTGAGCGG CGACGCCAAC GTGCGCATCC GCAACTCCCG CGAGGTGCTG	2220
TTCAAGAAGA GGTACATGAG CGGCGCCAAG GACGTGAGCG AGATGTTTAC CACCAAGTTC	2280
GAGAAGGACA ACTTCTACAT CGAGCTGAGC CAGGGCAACA ACCTGTACGG CGGCCCGATC	2340
GTGCACTTCT ACGACGTGAG CATCAAGTAG	2370

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2241 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA encoding VIP3A(c)"

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATGAACAAGA ACAACGCCAA GCTGAGCACC CGCGCCCTGC CGAGCTTCAT CGACTACTTC	60
AACGGCATCT ACGGCTTCGC CACCGGCATC AAGGACATCA TGAACATGAT CTTCAAGACC	120
GACACCGGCG GCGACCTGGC CCTGGACGAG ATCCTGGAGA ACCAGCAGCT GCTGAACGAC	180
ATCAGCGGCA AGCTGGACGG CGTGAACGGC AGCCTGAACG ACCTGATCGC CCAGGGCAAC	240
CTGAACACCG AGCTGAGCAA GGAGATCCTT AAGATCGCCA ACGAGCAGAA CCAGGTGCTG	300
AACGACGTGA ACAACAAGCT GGACGCCATC AACACCATGC TGC GCGTGTA CCTGCCGAAG	360
ATCACCAGCA TGCTGAGCGA CGTGATGAAG CAGAACTACG CCCTGAGCCT GCAGATCGAG	420
TACCTGAGCA AGCAGCTGCA GGAGATCAGC GACAAGCTGG ACATCATCAA CGTGAACGTC	480
CTGATCAACA GCACCCTGAC CGAGATCACC CCGGCCTACC AGCGCATCAA GTACGTGAAC	540
GAGAAGTTTCG AAGAGCTGAC CTTCGCCACC GAGACCAGCA GCAAGGTGAA GAAGGACGGC	600
AGCCCGGCGG ACATCCGGGA CGAGCTGAGC GAGCTGACCG AGCTGGCGAA GAGCGTGACC	660
CAGAACGACG TGGACGGCTT CGAGTTCTAC CTGAACACCT TCCACGACGT GATGGTGGGC	720
AACAACCTGT TCGGCCGAG CGCCCTGAAG ACCGCCAGCG AGCTGATCAC CAAGGAGAAC	780
GTGAAGACCA GCGGCAGCGA GGTGGGCAAC GTGTACAACT TCCTGATCGT GCTGACCGCC	840
CTGCAGGCC AGGCCTTCCT GACCCTGACC CCCTGTGCGA AGCTGCTGGG CCTGGCCGAC	900
ATCGACTACA CCAGCATCAT GAACGAGCAC TTGAACAAGG AGAAGGAGGA GTTCCGCGTG	960

AACATCCTGC CGACCCTGAG CAACACCTTC AGCAACCCGA ACTACGCCAA GGTGAAGGGC	1020
AGCGACGAGG ACGCCAAGAT GATCGTGGAG GCTAAGCCGG GCCACGCGTT GATCGGCTTC	1080
GAGATCAGCA ACGACAGCAT CACCGTGCTG AAGGTGTACG AGGCCAAGCT GAAGCAGAAC	1140
TACCAGGTGG ACAAGGACAG CTTGAGCGAG GTGATCTACG GCGACATGGA CAAGCTGCTG	1200
TGTCCGGACC AGAGCGGGCA AATCTACTAC ACCAACAACA TCGTGTTCCT GAACGAGTAC	1260
GTGATCACCA AGATCGACTT CACCAAGAAG ATGAAGACCC TCGCTACGA GGTGACCGCC	1320
AACTTCTACG ACAGCAGCAC CGGCGAGATC GACCTGAACA AGAAGAAGGT GGAGAGCAGC	1380
GAGGCCGAGT ACCGCACCCT GAGCGCGAAC GACGACGGCG TCTACATGCC ACTGGGCGTG	1440
ATCAGCGAGA CCTTCCTGAC CCCGATCAAC GGCTTTGGCC TGCAGGCCGA CGAGAACAGC	1500
CGCCTGATCA CCCTGACCTG TAAGAGCTAC CTGCGCGAGC TGCTGCTAGC CACCGACCTG	1560
AGCAACAAGG AGACCAAGCT GATCGTGCCA CCGAGCGGCT TCATCAGCAA CATCGTGGAG	1620
AACGGCAGCA TCGAGGAGGA CAACCTGGAG CCGTGAAGG CCAACAACAA GAACGCCTAC	1680
GTGGACCACA CCGCGGCGT GAACGGCACC AAGGCCCTGT ACGTGCACAA GGACGGCGGC	1740
ATCAGCCAGT TCATCGGCGA CAAGCTGAAG CCGAAGACCG AGTACGTGAT CCAGTACACC	1800
GTGAAGGGCA AGCCATCGAT TCACCTGAAG GACGAGAACA CCGGCTACAT CCACTACGAG	1860
GACACCAACA ACAACCTGGA GGACTACCAG ACCATCAACA AGCGCTTCAC CACCGGCACC	1920
GACCTGAAGG GCGGTACCT GATCCTGAAG AGCCAGAACG GCGACGAGGC CTGGGGCGAC	1980
AACTTCATCA TCCTGGAGAT CAGCCCGAGC GAGAAGCTGC TGAGCCCGGA GCTGATCAAC	2040
ACCAACAACCT GGACCAGCAC CGGCAGCACC AACATCAGCG GCAACACCCT GACCCTGTAC	2100
CAGGGCGGCC GCGGCATCCT GAAGCAGAAC CTGCAGCTGG ACAGCTTCAG CACCTACCGC	2160
GTGTACTTCA GCGTGAGCGG CGACGCCAAC GTGCGCATCC GCAACTCCCG CGAGGTGCTG	2220
TTCGAGAAGA AGGACAAGTA G	2241

What is claimed is:

1. A protein of the VIP3 class which is VIP3A(c) and homologues thereto.
- 2.. A protein wherein the amino acid sequence comprises a toxic domain of a protein of the VIP3 class.
3. The protein of claim 2 wherein the toxic domain is the toxic domain of a VIP3A(a) protein.
4. A transgenic plant comprising a DNA sequence which encodes a protein of the VIP3 class.
5. The transgenic plant of claim 4 wherein the DNA sequence encodes a VIP3A(a) protein.
6. The transgenic plant of claim 5 wherein the DNA sequence encodes a VIP3A(c) protein.
7. The transgenic plant of any one of claims 4 to 6 wherein said plant is selected from the group consisting of maize, sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape.
8. The transgenic plant of claim 7 wherein said plant is a maize plant.
9. The transgenic plant of any one of claims 4 to 8 further comprising a second DNA sequence which encodes a second insecticidal protein.
10. The transgenic plant of claim 9 wherein said second DNA sequence encodes a  $\delta$ -endotoxin, another protein of the VIP3 class, a protein of the VIP1 class or a protein of the VIP2 class.
11. The transgenic plant of claim 10 wherein said second DNA sequence is a  $\delta$ -endotoxin.
12. A microorganism comprising a heterologous DNA sequence wherein the DNA sequence encodes a VIP3A(a) protein.
13. The microorganism of claim 12 wherein the DNA sequence encodes a VIP3A(c) protein.
14. The microorganism of claims 12 or 13, wherein said microorganism is selected from the group consisting of bacteria, baculovirus, algae and fungi.
15. The microorganism of claim 14 wherein said microorganism is selected from the group consisting of *Bacillus*, *Pseudomonas*, *Clavibacter*, and *Rhizobium*.

16. The microorganism of any one of claims 12 to 15 further comprising a second DNA sequence which encodes a second insecticidal protein.
17. The microorganism of claim 16 wherein said second DNA sequence encodes a  $\delta$ -endotoxin, another protein of the VIP3 class, a protein of the VIP1 class or a protein of the VIP2 class.
18. The microorganism of claim 17 wherein said second DNA sequence is a  $\delta$ -endotoxin.
19. An entomocidal composition comprising microorganisms according to any one of claims 12 to 18.
20. A method of controlling insects by contacting the insects with an insecticidal amount of a chemical ligand to a receptor of the VIP3 class of proteins, or an antibody to a receptor of the VIP3 class of proteins.
21. The method of claim 20 wherein the insects are contacted with a transgenic plant comprising a DNA sequence which expresses a protein of the VIP3 class, preferably a VIP3A(c) protein.
22. The method of claim 20 wherein the insects are contacted with an entomocidal composition comprising a microorganism further comprising a heterologous DNA sequence capable of expressing a protein of the VIP3 class, preferably a VIP3A(c) protein.
23. A recombinant DNA sequence which encodes a VIP3A(c) protein and homologues thereto.
24. The recombinant DNA sequence of claim 23 wherein the DNA sequence is a synthetic sequence which has been designed for optimum expression in a plant.
25. The recombinant DNA sequence of claim 24 wherein the plant is a maize plant.
26. An expression cassette comprising a heterologous promoter operably linked to a DNA sequence encoding a protein of a VIP3A(c) protein.
27. The expression cassette of claim 26 wherein said promoter functions in plants is selected from the group consisting of inducible, constitutive, tissue-preferred and tissue-specific promoters.
28. The expression cassette of claim 27 wherein the promoter is selected from the group consisting of the ubiquitin, PEP carboxylase, LPT and MTL promoters.
29. A receptor to a protein of the VIP3 class.
30. A DNA sequence which encodes a receptor of the VIP3 class.
31. The receptor of claim 29 comprising a death domain and a repeated EGF-motif.

32. The receptor of claim 31 having the sequence set forth in SEQ ID NO:9.

33. The DNA sequence according to claim 30 as set forth in SEQ ID NO:8.

34. Antibodies to a receptor of the VIP3 class of proteins.

35. A method of identifying and isolating homologues of a receptor to a protein of the VIP3 class or of a DNA sequence which encodes a receptor to a protein of the VIP3 class comprising obtaining

(a) a DNA sequence which encodes a receptor to a protein of the VIP3 class, hybridizing said DNA sequence with DNA obtained from a test organism, detecting hybridization to the DNA from said organism, and isolating said homologue from said organism;

(b) a DNA sample from an organism, using primers to a DNA sequence encoding a receptor to a protein of the VIP3 class, obtaining a reaction product, then isolating a DNA sequence which encodes a receptor to a protein of the VIP3 class from said organism.

(c) a protein sample from a test organism, obtaining an antibody to a receptor to a protein of the VIP3 class, reacting said antibody with said protein sample, and detecting and isolating homologues by detecting the presence of an immunological reaction.

36 A method of identifying a compound as a VIP3 receptor chemical ligand having pesticidal activity comprising exposing the VIP3 receptor to a test compound, and assaying the interaction between the receptor and the test compound.

37. The method according to claim 36 wherein the VIP3 receptor is cellularly expressed and the assayed interaction is programmed cell death.

38. The method according to claim 36 wherein the assayed interaction is specific binding between the VIP3 receptor and the test compound.

Figure 1

1 20  
SGSPGLQEFAAASTMYSRIFLLVIVCAVKASLFTVN  
40 60  
VYDDNPETEIASSLKGCNPQECDQRCRRLKFPGGA  
80 100  
CVNGRCKCDNFLSVKDDVSVEEPAILKDLVSLEAEQ  
120 140  
AAKSRRCNRVCDAVCRALHNTSGACVDGQCKCTN  
160  
KISAGDIVSDPAESLRTCNPIRCDEQCRRNGHEFGV  
180 200  
CFKGQCKCDYFLKEEVDEPEVTSLPKNCNPQECDQ  
220 240  
RCRRLKFPGACVNGRCKCDNFFSAGDIVSDPAES  
260 280  
LRSCNPIRCDEQCRRNGHEFGVCFKGQCKCDYFL  
300  
NSEVDAVNEFPQAGSKRYCNLTQCNQTCANRFYD  
320 340  
SARVIHGWCKCYSKMERQDASPLNDVTEDENEVS  
360 380  
NDILRTVAEELSDVSPRACKSASCNQACRAFYFKG  
396  
GWCRFGRCQCF